

Recognition of Emotions in Autism: A Formal Meta-Analysis

Mirko Uljarevic · Antonia Hamilton

© Springer Science+Business Media New York 2012

Abstract Determining the integrity of emotion recognition in autistic spectrum disorder is important to our theoretical understanding of autism and to teaching social skills. Previous studies have reported both positive and negative results. Here, we take a formal meta-analytic approach, bringing together data from 48 papers testing over 980 participants with autism. Results show there is an emotion recognition difficulty in autism, with a mean effect size of 0.80 which reduces to 0.41 when a correction for publication bias is applied. Recognition of happiness was only marginally impaired in autism, but recognition of fear was marginally worse than recognition of happiness. This meta-analysis provides an opportunity to survey the state of emotion recognition research in autism and to outline potential future directions.

Keywords Autism · Emotion · Face · Meta-analysis · Social

Introduction

In Kanner's original (1943) description of autism, he considered this condition to be an "example of inborn

autistic disturbances of affective contact". Over 60 years later, the role of emotion in autism is still debated. Current ICD-10 and DSM-IV criteria for diagnosis of autistic spectrum condition (ASC) list marked impairments in the use of facial expression, body postures, and gestures to regulate social interaction; the lack of mutual sharing of emotions, impaired or deviant response to other people's emotions and the lack of spontaneous seeking to share enjoyment, among other symptoms. These difficulties in using, sharing and responding to emotions correspond roughly to two of the three components of emotion processing (as defined by Begeer et al. 2008; Herba and Phillips 2004; Phillips et al. 2003), namely production of an emotional state and regulation of that state. Diagnostic criteria for autism do not require a difficulty in the first of Phillips's components; the identification of emotional cues. Nevertheless, it is commonly assumed that emotion recognition difficulties are present in individuals with ASC.

In typical children, recognition of emotional facial expressions is an early developing social skill. Walker-Andrews (1998) found that 4-month-old infants were able to discriminate between expressions of anger, fear, sadness, happiness and surprise when those expressions were presented in a familiar context and that their reactions were specific for particular emotional expressions. Also, between 8 and 10 months infants begin to use emotional expressions for social referencing (Camras and Shutter 2010). Emotional expressions are a basic source of information about the sender's current emotional state (Ekman 1992), intentions (Adams et al. 2006) and about important objects and events in the environment (Moses et al. 2001; Olsson et al. 2007).

Failure of these fundamental early emotion recognition skills would have profound consequences for a child's social development, cutting the child off from learning

Electronic supplementary material The online version of this article (doi:10.1007/s10803-012-1695-5) contains supplementary material, which is available to authorized users.

M. Uljarevic
School of Psychology, Cardiff University, Cardiff, Wales, UK

A. Hamilton (✉)
School of Psychology, University of Nottingham, University
Park, Nottingham NG7 2RD, UK
e-mail: antonia.hamilton@nottingham.ac.uk

about other people's feelings and responses. Thus, it has been suggested that emotion reading might be a primary difficulty in autism (Hobson 1986a, b). Some interventions for autism specifically teach emotion recognition skills (Golan et al. 2010; Hopkins et al. 2011). However, despite numerous studies, there is not yet any consensus whether if basic emotion recognition is a fundamental and universal difficulty for individuals with ASC or not. The following brief review provides a summary of some of the research conducted (see Harms et al. 2010 for a more detailed approach).

Early work on emotion matching suggested that participants with autism have difficulty matching emotional facial expressions to emotional body actions, contexts or line drawings (Hobson 1986a, b; Hobson et al. 1988; Weeks and Hobson 1987; Braverman et al. 1989). However, a detailed study with well-matched participant groups did not find any evidence for basic emotion recognition difficulties (Ozonoff et al. 1990). Research turned to the idea that participants with autism might have difficulties in the recognition of just some of the six basic emotions rather than a generalised deficit. Baron-Cohen et al. (1993) suggested that theory of mind difficulties in autism could cause selective difficulties in recognizing surprise, and found some evidence in favour. However, other studies have failed to replicate this finding (Baron-Cohen et al. 1997; Castelli 2005; Spezio et al. 2007). Other studies suggest fear recognition is most difficult for individuals with autism (Ashwin et al. 2006; Corden et al. 2008; Howard et al. 2000; Humphreys et al. 2007; Pelphrey et al. 2002; Wallace et al. 2008). Difficulties have also been reported in other negative emotions (anger: Ashwin et al. 2006; disgust: Wallace et al. 2008; Humphreys et al. 2007; Ashwin et al. 2006; sadness: Boraston et al. 2007; Corden et al. 2008; Wallace et al. 2008). However, there are also published studies that did not find impairments in the recognition of fear and other negative emotions (Lacroix et al. 2009; Piggot et al. 2004) or found deficits in the recognition of positive emotions as well (Humphreys et al. 2007).

Research on generalised emotion recognition difficulties in ASC has also continued, with very mixed results. Published studies have found generalised deficits on various emotion reading tasks (Corbett et al. 2009; Davies et al. 1994; Loveland et al. 2008; Tantam et al. 1989). However, there are also a significant number of papers reporting no differences between typical and autistic participants (Baron-Cohen et al. 1997; Castelli 2005; Da Fonseca et al. 2009; Jones et al. 2011; Lacroix et al. 2009; Neumann et al. 2006; Ozonoff et al. 1990; Piggot et al. 2004; Spezio et al. 2007). These include some of the studies with the closest match between participant groups (Ozonoff et al. 1990) and studies with large sample sizes (Jones et al. 2011; Loveland et al. 2008).

Thus, the overall picture of emotion reading in autism is very mixed. The interpretation is further complicated by the large variability between studies in sample size, task, participant characteristics and group matching. Published studies have sample sizes ranging from only 5 participants up to 97, and it is possible that many studies are underpowered. Different tasks have also been used, but positive and negative results have been found in both emotion labelling tasks which might rely on verbal skills and in emotion matching tasks which are non-verbal. Some have suggested that subtle or difficult tasks are required to reveal emotion reading difficulties (Clark et al. 2008; Humphreys et al. 2007; Law Smith et al. 2010). Again, other studies show good performance by autistic children even in subtle tasks (Castelli 2005; Tracy et al. 2011). Finally, it has been suggested that deficits in emotion recognition are only evident when the autistic group is not carefully matched with the control group (Ozonoff et al. 1990). However, although this fact might explain findings on some earlier studies (Tantam et al. 1989), more recent research has addressed this issue more carefully (Humphreys et al. 2007; Wallace et al. 2008).

This brief narrative review demonstrates that there are currently no straightforward answers in research on emotion recognition in autism. It is not clear if individuals with autism spectrum disorder are impaired in their ability to read basic emotional expressions. Furthermore, if such impairment does exist, it is not clear if all emotions are equally affected or whether reading of certain emotions might be spared or impaired to a lesser extent. Ozonoff et al. (1990) suggested that for the emotion reading impairment to be considered a fundamental deficit in autism, impairments should be apparent across studies, paradigms and control groups. One way to test this is to conduct a formal meta-analysis.

Meta-analysis uses a strict set of search criteria to enable identification of all possible and relevant research studies published on the subject of interest. Inclusion and exclusion criteria are clearly stated and this together with comprehensive search minimizes the risk of bias. Statistical analysis of effect sizes for each study weighted by the sample size of the study provides numerical estimates of overall effect size, as well as the impact of moderator variables and the possibility of publication bias. By analyzing large collections of data from individual studies, meta-analysis can overcome the problem of heterogeneity in results due to small sample size and heterogeneity in study characteristics (Egger 1997a; Green 2005; Van den Noortgate and Onghena 2006). The present paper uses a formal meta-analytic approach to examine the question of emotion recognition in autism based on the existing literature. This approach allows us to systematically summarise and integrate the findings of multiple studies, and thus

assess the general question of emotion reading in autism. We focus on emotion reading from visual stimuli because these are most studied, and aim to determine if deficits in reading visual emotions are present across age, IQ and task in autism, and if such difficulties are equivalent in magnitude across different emotions.

Methods

Literature Search

In order to find eligible studies, we searched Web of Science, PsychINFO and PubMed using combinations of the following terms: autism, Asperger syndrome, pervasive developmental disorders, emotion recognition, emotion perception, facial expression, facial affect, face, body. Additional searches of relevant journals (Journal of Autism and Developmental Disorders, Autism, The Journal of Child Psychology and Psychiatry, Journal of Child Psychology and Psychiatry, Developmental Psychopathology) were performed. Also we searched the reference list of review articles and lists of publications of researchers working in this field. Studies published after December 2011 were not included.

Inclusion Criteria

We included studies published in English comparing a group of participants formally diagnosed with ASC and a group of typically developed subjects. Master and doctoral theses and conference presentations were not included. We limited our analysis to studies examining recognition of emotions presented in the visual modality. Information regarding the accuracy on behavioural tasks had to be available in order for study to be included. We included all studies examining more than one of the six standard emotions: fear, surprise, anger, disgust, happiness and surprise expressed by face and body (Ekman and Friesen 1976; Prinz 2004). Complex or social emotions and recognition of emotional hand gestures were excluded. Neuroimaging, electroencephalographic, eye-tracking and physiological studies were also included if they employed behavioural tasks that met above mentioned criteria. Figure 1 provides an overview of the selection process, and Table 3 of supplementary information lists the 59 articles which were excluded with details of why.

Coding and Analysis

Each study was initially coded by the first author and coding was then checked by the second author. Studies were coded for the following variables:




Fig. 1 Selection process. This chart indicates how papers were selected for inclusion in the meta-analysis. Articles were excluded if they were not relevant (e.g. did not examine participants with autism, did not examine recognition of emotion, were review articles, were not published in English)

1. Number/gender of participants in autistic and control groups;
2. Diagnosis and tools used for diagnosis;
3. Mean age and standard deviation of participants in autistic and control groups;
4. Level of function of participants—this included mental age (verbal and non verbal) intelligence (full scale, verbal and performance) and instruments used;
5. Type of task: we classified tasks as emotion labelling (EL) which could be forced choice or free choice, emotion matching (EM) or a different task (detailed in Table 1 of supplementary information).
6. Stimuli: whether stimuli showed faces or bodies, and whether they were static or dynamic. For face stimuli, the source and style of stimuli was noted.
7. Emotions: we focused on “the big six” (Ekman and Friesen 1976; Prinz 2004): happiness, anger, fear, disgust, sadness, surprise and not on the emotions later included in this category (amusement, embarrassment, excitement, pride, pride, shame, relief ...).

We coded the results of each study in terms of the mean correct performance of each group and the standard deviation of correct performance, rather than reaction time or neurophysiological measures. Reaction time data was not coded because many studies did not collect this data, but this does limit the sensitivity of our analysis. Where performance measures had been recorded but full data was not present in the paper (e.g. data was plotted but not listed as numbers), we contacted the authors to obtain the groups means and standard deviations. We are grateful to all authors who responded to our requests. Using the mean and SD data, we calculated Hedge’s *d*, a measure of effect size

that is equivalent to Cohen's *d* but includes a correction for small sample sizes (Hedges 1981). Studies that did not report means and standard deviations of behavioural performance for each group could not be included, but are listed in Table 3 of supplementary information. Where studies reported performance data for individual emotions separately, we recorded this data (Table 2 of supplementary information) to enable an analysis of differences in recognition performance for different emotions. Again, data was summarised in terms of Hedge's *d* and the variance of *d*. Supplementary data 4 lists the full references of all studies in the meta-analysis.

Some studies reported more than one analysis of the same ASC participant group. For example, the same participants might have completed an emotion-matching and an emotion-labelling task or might be matched to one typical group on verbal mental age and to another typical group on non-verbal mental age. Each of these results is listed as a separate row in Table 1 of supplementary information. However, it would not be appropriate to include data from exactly the same autism sample several times in the meta-analysis, so we had to select only one result from each study for further consideration. The comparisons which were included are marked with a star in Table 1 of supplementary information. Where studies tested the same group of participants on multiple tasks, we included the Emotion Labelling task (EL) in preference to others, and Static stimuli in preference to dynamic. This selection was made because these were the most common tasks/stimuli, and we wanted to reduce heterogeneity in our results. Analysis using the less common task/stimuli where possible did not change our results substantially. Where multiple comparison groups were present in a study, we included the group matched for verbal abilities in preference to a non-verbal match.

We conducted three analyses on the effect size data. First, analysis of effect size and assessment of the role of moderator variables (participant age and IQ) was conducted using MetaWin 2.0 (Rosenberg et al. 2007). This allows us to answer the general question of whether participants with autism do show emotion recognition difficulties and to obtain confidence limits on this answer. Second, analysis of possible publication bias and the implementation of the trim-and-fill method for ameliorating publication bias (Duval and Tweedie 2000) was implemented using the Meta package in R (Schwarzer 2007; <http://cran.r-project.org/web/packages/meta/index.html>). This allows an initial assessment of whether unpublished data on emotion recognition (likely to have null results) is an important issue in this area. Finally, we examined effect sizes in the 16 studies which reported data from individual emotions. Here we used conventional ANOVA and *t* tests in SPSS to determine if there were difference in effect size between different emotions.

Results

48 papers published from 1989 to 2011, which tested 932 participants with autism met our criteria (see Fig. 1 for the description of selection process and Table 3 of supplementary information for the list of excluded studies). Table 1 of supplementary information presents a list of the studies that were included in our analysis. Twenty eight comparisons used the Ekman facial affect set of stimuli and 46 used different stimuli. 65 comparisons used static faces and 10 used dynamic stimuli. 38 comparisons used an emotion labelling task (32 forced and 6 free labelling task), 23 used an emotional face matching task and the remaining 12 used other tasks. 63 comparisons examined recognition of facial emotions, while 7 comparisons used body emotion and 8 used emotional context tasks. Sample sizes ranged from 5 (Pelphrey et al. 2002) to 97 (Jones et al. 2011). Mean age of participant's ranged from 6 to 41 years, and mean FSIQs ranged from 40 to 130. Thus, our meta-analysis covers the full range of ages, IQs and emotion recognition tasks which are encountered in the research literature.

General Emotion-Recognition Difficulties in ASC

For this analysis, we included 50 comparisons that examined emotion recognition without overlap in the participant groups. From a random effects analysis of overall effect size, we found a mean effect size of -0.800 , with 95 % confidence limits from -0.57 to -0.99 . Negative effect sizes indicate worse performance by the autism group. This is a large effect size (Cohen 1988) and indicates that overall, individuals with autism do have difficulties in emotion recognition. The studies also had substantial heterogeneity ($Q_{total} = 77.83$, $df = 49$, $p = 0.0054$), indicating that effect sizes were not uniform across studies (Gurevitch and Hedges 1999).

We considered the impact of two moderator variables on emotion recognition—the mean age of the autism sample and the mean IQ of the autism sample. Two outlier studies with very large negative effect sizes which were more than 2 standard deviations from the mean effect size (visible on the left of Fig. 2; Da Fonseca et al. 2009; Gepner et al. 1996) were excluded from this analysis. There were no significant effects of age (slope = -0.01 , $p = 0.24$) or of IQ (slope = -0.004 , $p = 0.27$) on effect size.

To examine effects of task, we compared 34 studies which used an emotion-labelling task to 13 studies which used an emotion matching task. Again, the two outlier studies were excluded. Effect size was -0.68 for emotion-matching and -0.70 for emotion labelling, with no difference between tasks ($Q = 0.007$, $df = 1,46$, $p = 0.93$). Thus, there was no evidence for systematic effects of task.




Fig. 2 Funnel Plot. Effect size for each study (Cohen's d) is plotted against the number of participants with autism in that study. *Filled circles* indicate studies in the meta-analysis. *Open squares* indicate studies inferred in the trim-and-fill analysis

The Impact of Publication Bias

Publication bias, arising when studies with null findings are not published, is an important issue to consider in any meta-analysis. As an initial test of publication bias, we plotted effect sizes against the number of autistic participants in a funnel plot (Egger et al. 1997b) (Fig. 2, solid circles). The clear asymmetry in the funnel plot, with the majority of studies to the left of the plot, suggests that publication bias may be an issue. Thus, we implemented Duval & Tweedie's trim-and-fill method, which estimated an additional 13 'missing studies' (open squares in Fig. 2). Repeating our standard random effects analysis with these filled studies gave an estimated effect size of -0.414 , with 95 % confident limits from -0.646 to -0.182 and a p value of 0.0005. Thus even after trim-and-fill, the overall effect size for the recognition of emotions by participants with autism is negative with confidence limits that do not span zero. It is important to note that the magnitude of the effect after trim-and-fill is substantially smaller than the raw estimate. Figure 1 also illustrates how the majority of published studies have small participant groups, with only 13 studies out of 45 testing 20 or more participants, and that the largest study reports an effect size close to zero.

Differences Between Individual Emotions

16 studies were available with data on the recognition of different emotions in individuals with autism (see Table 2 of supplementary information). These tested 379 participants with autism and most examined adults. Three of these studies used more than one task on the same participant sample, so all results are listed but only one comparison from each study was included in the meta-analysis. All six individual emotions showed negative effect sizes. For five emotions (sadness, anger, surprise, fear, disgust),

the 95 % confidence intervals were entirely in the negative range, suggesting that adults with ASC have difficulty with recognition of each of these individual emotions. However, the 95 % confidence intervals for happiness spanned zero, demonstrating no reliable difficulty in the recognition of happiness across these studies (Fig. 2). As the confidence limits came close to zero, it is possible that there is a marginal difficulty in the recognition of happiness which would be revealed with more subtle measures such as reaction time.

Comparison between different emotions was complicated by the fact that only eight of these studies tested all six emotions (Table 2 of supplementary information). An ANOVA across the effect sizes found in these eight studies revealed no significant effects of emotion ($F = 1.89$; $df = 5,35$; $p = 0.12$). This is congruent with Fig. 3 which shows overlapping confidence intervals between all emotions. To allow all the original data to be considered, we decided to compare each emotion to Happiness as a baseline emotion. Unfortunately, there was not enough data on recognition of neutral faces (the more traditional baseline) to use this option. Happiness makes a suitable baseline because no theoretical accounts suggest a specific impairment in happiness, and because all studies tested this item. We performed paired-sample t tests comparing the effect size for each other emotion to the effect size for happiness. Results revealed no difference for sadness ($p = 0.36$, $df = 15$, $t = 0.94$), surprise ($p = 0.14$, $df = 10$, $t = 1.6$), disgust ($p = 0.32$, $df = 9$, $t = 1.06$) or anger ($p = 0.069$, $df = 14$, $t = 1.97$). Recognition of fear ($p = 0.044$, $df = 12$, $t = 2.248$) was significantly worse than recognition of happiness, but as this result would not survive Bonferroni correction for 5 comparisons ($p < 0.01$), it must be considered marginal.

Discussion

This paper used formal meta-analysis to examine whether individuals with autism spectrum disorders show general emotion recognition deficits. We examined 48 studies testing over 930 participants with autism, and found evidence of a large negative effect size (-0.80) indicating that there is indeed a general impairment in emotion recognition in individuals with ASC. This effect size was substantially reduced (to -0.41) but still significantly different from zero when a correction for publication bias was included. Participant age, IQ and task had no impact on performance. There was marginal evidence for differences between emotions, with confidence limits for recognition of happiness spanning zero, and marginally worse recognition of fear than happiness. Based on these results, we discuss general methodological issues in emotion research




Fig. 3 Effect size for different emotions. Mean effect size ($\pm 95\%$ confidence limits) are plotted for each of the six basic emotions. Data are from Table 2 of supplementary information

and then consider the impact of age, IQ, task and the different emotions tested on performance by participants with autism. Finally, we highlight useful directions for further research.

Methodological Issues

The question of how best to examine atypical development of cognitive abilities is not an easy one to answer (Charman et al. 1998; Johnson et al. 2002; Thomas et al. 2009) and our overview of the literature raises several important points. First, the issue of publication bias (Dickersin 1990) is likely to be important in autism-emotion research. Publication of null effects tends to be harder than publication of positive results, and it is plausible that studies with small groups and null findings (or better performance in the autism group) remain hidden in university filing cabinets around the world. Even in cases where results are published, statistical tests that did not yield significant results (at $p < 0.05$) are sometimes not reported in detail. Of the 58 comparisons in Table 3 of supplementary information (which did not provide detailed results), 22 reported either no differences, mixed results or subtle differences between typical and autistic groups. In contrast, the studies in Table 1 of supplementary information (which did provide detailed results) reported significant findings in the vast majority of cases. This suggests a bias in reporting statistics even within published papers, and motivates us to strongly encourage researchers to report in full the results of the statistical tests they conducted, even when not significant.

Second, of the 48 studies in Table 1 of supplementary information, only 31% (15 studies) tested groups with 20 or more participants. However, a straightforward power analysis suggests that detection of a large group difference (effect size = 0.8) between two independent populations with a power of 0.95 requires 35 participants in each group. This suggests that the vast majority of studies examining

emotion recognition in autism are underpowered. This is where a meta-analysis of the form conducted here can add substantially to the existing literature. Notably, the two largest studies in our sample (Jones et al. 2011; Loveland et al. 2008), which examined 97 and 80 autistic participants, respectively, both found no evidence of group differences in emotion recognition.

We used Duval & Tweedie's trim-and-fill method to attempt to correct for publication bias, and this analysis 'filled in' an additional 13 results. After filling-in, the overall effect size for the emotion recognition deficit in autism remained significantly different from zero but was small (0.40). Power analysis suggests that groups of over 135 participants would be needed to reliably detect such an effect in a simple comparison of typical and autistic participants, but no researchers have yet attempted such an ambitious task. However, we note a trend for larger sample sizes in more recent studies and hope this continues.

Age, IQ and Task Factors

Our meta-analysis shows that the question of emotion recognition in autism is an active one which continues to attract research 25 years after the first studies on the topic. With over 100 articles published on the topic of emotion processing in autism, it might seem surprising that this question still remains unresolved. One possible explanation for this might be the heterogeneity of tasks and participant groups involved in different research studies. Our studies covered a wide age range but did not reveal any effects of age or IQ on emotion recognition performance. This suggests that emotion recognition difficulties are not specific to any particular subgroup of individuals with autism (e.g. lower functioning individuals), and that there were no substantial changes in recognition performance with age. This does not mean that individuals cannot improve as they grow older, but rather that the population as a whole does not improve. We also note that this lack of an IQ effect does not mean that there is no relationship between IQ and emotion processing. Many studies in the meta-analysis match participant groups on the basis of their IQ, so these studies can at best show participants with ASC performing at the level expected for their mental age, not at the same level as individuals with the same chronological age. Second, several studies in the meta-analysis provide only very limited IQ information. Thus, further research would be needed to define how intellectual capacity relates to emotion processing.

Another important issue in considering the role of IQ and level of functioning in emotion processing is the possibility of subgroups within the autism spectrum having different capabilities. The autism spectrum can be divided into classical autism, high-functioning autism, Asperger's

syndrome and pervasive developmental disorder not otherwise specified (PDD-NOS) though the meaning of these categories is under debate (Lord and Jones 2012; Mandy et al. 2012). It is possible that emotion processing is abnormal in just some of these groups but not others. Unfortunately, the papers reviewed in this meta-analysis used a variety of diagnostic tools and few distinguish between these different subgroups of autism. Thus, it was not feasible to examine emotion recognition within specific subgroups. This would be an interesting topic for future studies.

Differences in the tasks used to assess emotion recognition could also contribute to the heterogeneity of results. In the studies we examined, 38 comparisons used an emotion labelling while 23 used an emotion matching task and 12 used other tasks. In a typical matching task, participants are asked to match pictures with a target picture where correct choice expresses the same emotion as the target but differs in other features such as identity or angle of view. In emotion labelling tasks participants are shown images or videos of emotional displays and are asked to either choose an item from a short, pre-specified list (forced choice format) or to come up with an emotion term (free labelling format) for what is portrayed on the picture.

While it is often assumed that both labelling and matching tasks tap the same core emotion recognition systems, there are important differences between them (Hariri et al. 2000; Herba and Phillips 2004; Phan et al. 2002). Matching tasks could be completed based on surface characteristics of the stimuli without a full understanding of the emotion. Thus, these tasks might lack sensitivity and allow individuals with autism to use compensatory strategies (Celani et al. 1999; Fein et al. 1992; Klin et al. 2002; Teunisse and de Gelder 2001). Labelling tasks require good verbal skills, especially in free-labelling conditions, but forced choice labelling may allow participants to guess a correct answer (Russell et al. 2003). Our meta-analysis did not find any evidence for overall differences in performance between emotion labelling and emotion matching tasks. This suggests that the difficulties experienced by ASC participants in these tasks are due to emotion processing and not to the linguistic or perceptual demands of these different tasks.

Role of Different Emotions

We analysed recognition of individual emotions in 16 studies that provided sufficient data and found that ASC individuals had difficulties in the recognition of five basic emotions but did not have difficulties in recognition of happiness (there may be a marginal difficulty here because confidence limits only just span zero). There was tentative evidence for worse recognition of fear than of happiness,

but no differences between happiness and sadness, surprise or disgust. One limitation of these results is that there was not enough data on the recognition of neutral faces for this category to contribute to our analysis. This means that we lack the ideal baseline, and had to use happiness as a 'baseline' emotion because this was the only emotion tested in all the studies examined. With these caveats, there are two important implications to these results. First, if recognition of happiness is not impaired in autism, this argues against the idea that poor emotion recognition is universal and primary in autism. Second, if recognition of fear is worse than recognition of happiness, this favours theories that link autism to poor eye contact and poor fear processing in the amygdala. We consider each of these in turn.

The finding that recognition of happiness is only borderline-impaired in autism might seem contrary to the global meta-analysis which found an overall recognition deficit. Mean effect size for happiness recognition was negative and the confidence limits only just spanned zero, suggesting there might be a marginal difference. Thus, it might be tempting to argue that more studies would reveal a true happiness recognition deficit. However, if publication bias is a factor, and our analysis above suggests this is likely, then studies reporting group differences in emotion recognition are more likely to be published than those which do not. This means that current estimates of effect size may be inflated, and future studies might decrease our estimate of effect size in happiness recognition and solidify the conclusion that recognition of this emotion is intact in autism. This is an important result because it suggests we should rule out theories that claim a global emotion recognition difficulty is primary and universal in autism. Rather, difficulties with emotion processing must be specific to particular emotions or stimuli.

A hint of specific difficulties was seen in the comparison of happiness recognition to fear recognition, where a marginally significant difference was found. Several theories link predict poor fear processing in autism, drawing on neurological or behavioural explanations. In neurological terms, it has been suggested that the amygdala has a specific role in the processing of fear (Adolphs 2008) and negative emotions in general (Adolphs et al. 1999; Anderson et al. 2000). Dysfunction of the amygdala in autism could cause poor recognition of fear and other negative emotions (Ashwin et al. 2006; Baron-Cohen et al. 2000; Howard et al. 2000), which is compatible with our data. Dysfunction of the amygdala in autism might lead to a lack of orienting to social stimuli, in particular to the eyes in a face (Neumann et al. 2006; Spezio et al. 2007). For example, several studies have found reduced attention to the eyes (Boraston et al. 2007; Dalton et al. 2005; Klin et al. 2002; Pelphrey et al. 2002) and increased attention to

the mouth region (Joseph and Tanaka 2003) in ASC, though contradictory results have been reported (Lopez et al. 2004; Van Der Geest et al. 2002). Processing of the eye region is particularly relevant to the recognition of fear, which requires attention to eyes and eye-brows (Dimberg and Petterson 2000; Dimberg and Thunberg 1998; Ekman 2004; Smith et al. 2005). In contrast, processing of the mouth region could be sufficient to judge happiness, which seems easier for participants with autism. Thus, amygdala dysfunction in autism could lead to reduced fixation on the eyes and to a difficulty in fear and anger recognition (Adolphs et al. 2005) together with better happiness recognition.

While this explanation is appealing, it is complicated by some results which suggest no difference between typical and autistic amygdala activity during emotion labelling and matching tasks (Piggot et al. 2004). There is also evidence that the amygdala does not respond only to fear, but functions as a ‘motivational relevance detector’ (Whalen 2007), which responds to positive and ambiguous stimuli as well (Phan et al. 2002; Whalen 2007). Thus, the status of an amygdala explanation of poor fear recognition in autism remains unclear.

Finally, our data provide evidence against one particular emotion-specific account of autism. At least some formulations of the Theory of Mind hypothesis predict a specific deficit in recognition of surprise in autism (Baron-Cohen et al. 1993). Of the six basic emotions, surprise is the only one that requires assessment of another person’s mental state (he expected something different, he is surprised). This means that if mental state judgements are impaired in autism and are required for processing of surprised facial expressions, then recognition of surprise might be specifically impaired. However, our results did not provide any evidence that surprise recognition is more difficult than recognition of any other emotions.

Future Directions

Ozonoff et al. (1990) suggested that for the emotion recognition difficulties to be considered a fundamental deficit in autism, impairments should be apparent across studies, paradigms and control groups. The data reviewed here do not strongly support a global emotion recognition impairment, because recognition of happiness was (just) intact across the studies we sampled. However, even a meta-analysis of this scale is fundamentally limited by the quality of the input data. Thus, issues such as sample size, group matching and the tasks used are critical (Burack et al. 2004; Harms et al. 2010; Jarrold and Brock 2004; Mervis and Klein-Tasman 2004). Based on the studies reviewed, we would encourage researchers in the field to use larger sample sizes in order to increase the reliability

and replicability of data. We would also strongly encourage full reporting of results (in tables, not just graphs) and of all statistical tests, even those which were not formally significant. Lack of full data substantially reduced the sample of published papers which could contribute to this meta-analysis (Table 3 of supplementary information).

Despite over 20 years of research, the status of emotion recognition in autism remains uncertain and the present meta-analysis highlights some possibilities. One important question is the role of timing in emotion recognition—individuals with autism might be slower to recognise emotions, or might have more difficulty with dynamically moving faces which have higher ecological validity than static photos. Examining emotion recognition in dynamic, time constrained and realistic contexts will be an important focus of future research. A second key area to focus on is potential differences in the recognition of different emotions, which has both theoretical and practical implications. Our results provide tentative evidence for poorer recognition of negative emotion, but further work testing different emotions in large participant groups and in combination with neuroimaging and eye tracking methods would be valuable. In particular, it is critical to determine the contribution of specific brain regions and abnormal eye scanning patterns to differences in recognition of different emotions. Addressing these questions will require more ambitious and large scale studies than cognitive scientists are accustomed to, but will provide critical insights into the origins of poor social cognition in autism, and the relationship between brain, development and social information processing.

Acknowledgments We thank Uta Frith, Tony Atkinson and Anneli Kylliainen for helpful comments on this manuscript, and Zoran Uljarevic for help preparing parts of the tables.

References

- Adams, R., Ambady, N., Macrae, C., & Kleck, R. (2006). *Emotional expressions forecast approach-avoidance behavior, motivation and emotion* (pp. 177–186). The Netherlands: Springer.
- Adolphs, R. (2008). Fear, faces, and the human amygdala. *Current Opinion in Neurobiology*, 18, 166–172.
- Adolphs, R., Gosselin, F., Buchanan, T. W., Tranel, D., Schyns, P., & Damasio, A. R. (2005). A mechanism for impaired fear recognition after amygdala damage. *Nature*, 433, 68–72.
- Adolphs, R., Tranel, D., Hamann, S., Young, A. W., Calder, A. J., Phelps, E. A., et al. (1999). Recognition of facial emotion in nine individuals with bilateral amygdala damage. *Neuropsychologia*, 37, 1111–1117.
- Anderson, A. K., Spencer, D. D., Fulbright, R. K., & Phelps, E. A. (2000). Contribution of the anteromedial temporal lobes to the evaluation of facial emotion. *Neuropsychology*, 14, 526–536.
- Ashwin, C., Chapman, E., Colle, L., & Baron-Cohen, S. (2006). Impaired recognition of negative basic emotions in autism: A test of the amygdala theory. *Social Neuroscience*, 1, 349–363.

- Baron-Cohen, S., Jolliffe, T., Mortimore, C., & Robertson, M. (1997). Another advanced test of theory of mind: Evidence from very high functioning adults with autism or Asperger syndrome. *Journal of Child Psychology and Psychiatry*, *38*, 813–822.
- Baron-Cohen, S., Ring, H. A., Bullmore, E. T., Wheelwright, S., Ashwin, C., & Williams, S. C. R. (2000). The amygdala theory of autism. *Neuroscience and Biobehavioral Reviews*, *24*, 355–364.
- Baron-Cohen, S., Spitz, A., & Cross, P. (1993). Do children with autism recognise surprise? A research note. *Cognition and Emotion*, *7*, 507–516.
- Begeer, S., Koot, H. M., Rieffe, C., Meerum Terwogt, M., & Stegge, H. (2008). Emotional competence in children with autism: Diagnostic criteria and empirical evidence. *Developmental Review*, *28*, 342–369.
- Boraston, Z., Blakemore, S. J., Chilvers, R., & Skuse, D. (2007). Impaired sadness recognition is linked to social interaction deficit in autism. *Neuropsychologia*, *45*, 1501–1510.
- Braverman, M., Fein, D., Lucci, D., & Waterhouse, L. (1989). Affect comprehension in children with pervasive developmental disorders. *Journal of Autism and Developmental Disorders*, *19*, 301–316.
- Burack, J., Iarocci, G., Flanagan, T., & Bowler, D. (2004). On mosaics and melting pots: Conceptual considerations of comparison and matching strategies. *Journal of Autism and Developmental Disorders*, *34*, 65–73.
- Camras, L. A., & Shutter, J. M. (2010). Emotional facial expressions in infancy. *Emotion Review*, *2*, 120–129.
- Castelli, F. (2005). Understanding emotions from standardized facial expressions in autism and normal development. *Autism*, *9*, 428–449.
- Celani, G., Battacchi, M. W., & Arcidiacono, L. (1999). The understanding of the emotional meaning of facial expressions in people with autism. *Journal of Autism and Developmental Disorders*, *29*, 57–66.
- Charman, T., Swettenham, J., Baron-Cohen, S., Cox, A., Baird, G., & Drew, A. (1998). An experimental investigation of social-cognitive abilities in infants with autism: Clinical implications. *Infant Mental Health Journal*, *19*, 260–275.
- Clark, T. F., Winkielman, P., & McIntosh, D. N. (2008). Autism and the extraction of emotion from briefly presented facial expressions: Stumbling at the first step of empathy. *Emotion*, *8*, 803–809.
- Cohen, J. (1988). *Statistical power analysis for the behavioral sciences (2nd edn)*. Academic Press: Routledge.
- Corbett, B. A., Carmean, V., Ravizza, S., Wendelken, C., Henry, M. L., Carter, C., et al. (2009). A functional and structural study of emotion and face processing in children with autism. *Psychiatry Research: Neuroimaging*, *173*, 196–205.
- Corden, B., Chilvers, R., & Skuse, D. (2008). Avoidance of emotionally arousing stimuli predicts social-perceptual impairment in Asperger's syndrome. *Neuropsychologia*, *46*, 137–147.
- Da Fonseca, D., Santos, A., Bastard-Rosset, D., Rondan, C., Poinso, F., & Deruelle, C. (2009). Can children with autistic spectrum disorders extract emotions out of contextual cues? *Research in Autism Spectrum Disorders*, *3*, 50–56.
- Dalton, K. M., Nacewicz, B. M., Johnstone, T., Schaefer, H. S., Gernsbacher, M. A., Goldsmith, H. H., et al. (2005). Gaze fixation and the neural circuitry of face processing in autism. *Nature*, *8*, 519–526.
- Davies, S., Bishop, D., Manstead, A. S. R., & Tantam, D. (1994). Face perception in children with autism and Asperger's syndrome. *Journal of Child Psychology and Psychiatry*, *35*, 1033–1057.
- Dickersin, K. (1990). The existence of publication bias and risk factors for its occurrence. *JAMA*, *263*, 1385–1389.
- Dimberg, U., & Petterson, M. (2000). Facial reactions to happy and angry facial expressions: Evidence for right hemisphere dominance. *Psychophysiology*, *37*, 693–696.
- Dimberg, U., & Thunberg, M. (1998). Rapid facial reactions to emotional facial expressions. *Scandinavian Journal of Psychology*, *39*, 39–45.
- Duval, S., & Tweedie, R. (2000). A nonparametric “Trim and Fill” method of accounting for publication bias in meta-analysis. *Journal of the American Statistical Association*, *95*, 89–98.
- Egger, M., Smith, G. D., & Phillips, A. N. (1997a). Meta-analysis: Principles and procedures. *BMJ*, *315*, 1533–1537.
- Egger, M., Smith, G. D., Schneider, M., & Minder, C. (1997b). Bias in meta-analysis detected by a simple, graphical test. *BMJ*, *315*, 629–634.
- Ekman, P. (1992). Facial expressions of emotion: An old controversy and new findings. *Philosophical Transactions of the Royal Society of London. Series B, Biological sciences*, *335*, 63–69.
- Ekman, P. (2004). *Emotions revealed: Recognizing faces and feelings to improve communication and emotional life*. New York: Owl Books.
- Ekman, P., & Friesen, W. V. (1976). Measuring facial movement. *Journal of Nonverbal Behavior*, *1*, 56–75.
- Fein, D., Lucci, D., Braverman, M., & Waterhouse, L. (1992). Comprehension of affect in context in children with pervasive developmental disorders. *Journal of Child Psychology and Psychiatry*, *33*, 1157–1162.
- Gepner, B., Gelder, B. D., & Schonen, S. D. (1996). Face processing in autistics: Evidence for a generalised deficit? *Child Neuropsychology*, *2*, 123–139.
- Golan, O., Ashwin, E., Granader, Y., McClintock, S., Day, K., Leggett, V., et al. (2010). Enhancing emotion recognition in children with autism spectrum conditions: An intervention using animated vehicles with real emotional faces. *Journal of Autism and Developmental Disorders*, *40*, 269–279.
- Green, S. (2005). Systematic reviews and meta-analysis. *Singapore Medical Journal*, *46*, 270–273. (quiz 274).
- Gurevitch, J., & Hedges, L. V. (1999). Statistical issues in ecological meta-analyses. *Ecology*, *80*, 1142–1149.
- Hariri, A. R., Bookheimer, S. Y., & Mazziotta, J. C. (2000). Modulating emotional responses: Effects of a neocortical network on the limbic system. *NeuroReport*, *11*, 43–48.
- Harms, M., Martin, A., & Wallace, G. (2010). Facial emotion recognition in autism spectrum disorders: A review of behavioral and neuroimaging studies. *Neuropsychology Review*, *20*, 290–322.
- Hedges, L. V. (1981). Distribution theory for glass's estimator of effect size and related estimators. *Journal of Educational and Behavioral Statistics*, *6*, 107–128.
- Herba, C., & Phillips, M. (2004). Development of facial expression recognition from childhood to adolescence: Behavioural and neurological perspectives. *Journal of Child Psychology and Psychiatry*, *45*, 1185–1198.
- Hobson, R. P. (1986a). The autistic child's appraisal of expressions of emotion. *Journal of Child Psychology and Psychiatry*, *27*, 321–342.
- Hobson, R. P. (1986b). The autistic child's appraisal of expressions of emotion: A further study. *Journal of Child Psychology and Psychiatry*, *27*, 671–680.
- Hobson, R. P., Ouston, J., & Lee, A. (1988). Emotion recognition in autism: Coordinating faces and voices. *Psychological Medicine*, *18*, 911–923.
- Hopkins, I. M., Gower, M. W., Perez, T. A., Smith, D. S., Amthor, F. R., Wimsatt, F. C., & Biasini, F. J. (2011). Avatar assistant: Improving social skills in students with an ASC through a computer-based intervention. *Journal of Autism Developmental Disorder*, *41*(11), 1543–1555.

- Howard, M. A., Cowell, P. E., Boucher, J., Brooks, P., Mayes, A., Farrant, A., et al. (2000). Convergent neuroanatomical and behavioural evidence of an amygdala hypothesis of autism. *NeuroReport*, *11*, 2931–2935.
- Humphreys, K., Minshew, N., Leonard, G. L., & Behrmann, M. (2007). A fine-grained analysis of facial expression processing in high-functioning adults with autism. *Neuropsychologia*, *45*, 685–695.
- Jarrold, C., & Brock, J. (2004). To match or not to match? Methodological issues in autism-related research. *Journal of Autism and Developmental Disorders*, *34*, 81–86.
- Johnson, M. H., Halit, H., Grice, S. J., & Karmiloff-Smith, A. (2002). Neuroimaging of typical and atypical development: A perspective from multiple levels of analysis. *Development and Psychopathology*, *14*, 521–536.
- Jones, C. R. G., Pickles, A., Falcaro, M., Marsden, A. J. S., Happe, F., Scott, S. K., et al. (2011). A multimodal approach to emotion recognition ability in autism spectrum disorders. *Journal of Child Psychology and Psychiatry*, *52*, 275–285.
- Joseph, R. M., & Tanaka, J. (2003). Holistic and part-based face recognition in children with autism. *Journal of Child Psychology and Psychiatry*, *44*, 529–542.
- Kanner, L. (1943). Autistic disturbances of affective contact. *Nervous Child*, *2*, 217–250.
- Klin, A., Jones, W., Schultz, R., Volkmar, F., & Cohen, D. (2002). Visual fixation patterns during viewing of naturalistic social situations as predictors of social competence in individuals with autism. *Archives of General Psychiatry*, *59*, 809–816.
- Lacroix, A., Guidetti, M., Rogé, B., & Reilly, J. (2009). Recognition of emotional and nonemotional facial expressions: A comparison between Williams syndrome and autism. *Research in Developmental Disabilities*, *30*, 976–985.
- Law Smith, M. J., Montagne, B., Perrett, D. I., Gill, M., & Gallagher, L. (2010). Detecting subtle facial emotion recognition deficits in high-functioning autism using dynamic stimuli of varying intensities. *Neuropsychologia*, *48*, 2777–2781.
- Lopez, B., Donnelly, N., Hadwin, J., & Leekam, S. (2004). Face processing in high-functioning adolescents with autism: Evidence for weak central coherence. *Visual Cognition*, *11*, 673–688.
- Lord, C., & Jones, R. M. (2012). Annual research review: Re-thinking the classification of autism spectrum disorders. *Journal of Child Psychology and Psychiatry*, *53*(5), 490–509.
- Loveland, K. A., Bachevalier, J., Pearson, D. A., & Lane, D. M. (2008). Fronto-limbic functioning in children and adolescents with and without autism. *Neuropsychologia*, *46*, 49–62.
- Mandy, W. P. L., Charman, T., & Skuse, D. H. (2012). Testing the construct validity of proposed criteria for DSM-5 autism spectrum disorder. *Journal of the American Academy of Child and Adolescent Psychiatry*, *51*(1), 41–50.
- Mervis, C. B., & Klein-Tasman, B. P. (2004). Methodological issues in group-matching designs: Alpha levels for control variable comparisons and measurement characteristics of control and target variables: Research methodology-matching. *Journal of Autism and Developmental Disorders*, *34*, 7–17.
- Moses, L. J., Baldwin, D. A., Rosicky, J. G., & Tidball, G. (2001). Evidence for referential understanding in the emotions domain at twelve and eighteen months. *Child Development*, *72*, 718–735.
- Neumann, D., Spezio, M. L., Piven, J., & Adolphs, R. (2006). Looking you in the mouth: Abnormal gaze in autism resulting from impaired top-down modulation of visual attention. *Social Cognitive and Affective Neuroscience*, *1*, 194–202.
- Olsson, A., Nearing, K. I., & Phelps, E. A. (2007). Learning fears by observing others: The neural systems of social fear transmission. *Social Cognitive and Affective Neuroscience*, *2*, 3–11.
- Ozonoff, S., Pennington, B., & Rogers, S. (1990). Are there emotion perception deficits in young autistic children? *Journal of Child Psychology and Psychiatry and Allied Disciplines*, *31*, 343–361.
- Pelphrey, K. A., Sasson, N. J., Reznick, J. S., Paul, G., Goldman, B. D., & Piven, J. (2002). Visual Scanning of Faces in Autism. *Journal of Autism and Developmental Disorders*, *32*, 249–261.
- Phan, K. L., Wager, T., Taylor, S. F., & Liberzon, I. (2002). Functional neuroanatomy of emotion: A meta-analysis of emotion activation studies in PET and fMRI. *NeuroImage*, *16*, 331–348.
- Phillips, M. L., Drevets, W. C., Rauch, S. L., & Lane, R. (2003). Neurobiology of emotion perception I: The neural basis of normal emotion perception. *Biological Psychiatry*, *54*, 504–514.
- Piggot, J., Kwon, H., Mobbs, D., Blasey, C., Lotspeich, L., Menon, V., et al. (2004). Emotional attribution in high-functioning individuals with autistic spectrum disorder: A functional imaging study. *Journal of the American Academy of Child and Adolescent Psychiatry*, *43*, 473–480.
- Prinz, J. J. (2004). Which emotions are basic? In D. Evans & P. Cruse (Eds.), *Emotion, evolution and rationality*. Oxford: Oxford University Press.
- Rosenberg, M. S., Adams, D. C., & Gurevitch, J. (2007). *MetaWin: Statistical software for meta-analysis 2.0*.
- Russell, J. A., Bachorowski, J.-A., & Fernandez-Dols, J.-M. (2003). Facial and vocal expressions of emotion. *Annual Review of Psychology*, *54*, 329–349.
- Schwarzer, G. (2007). Meta: An R package for meta-analysis. *R News*, *7*(3), 40–45.
- Smith, M. L., Cottrell, G. W., Gosselin, F., & Schyns, P. G. (2005). Transmitting and decoding facial expressions. *Psychological Science*, *16*, 184–189.
- Spezio, M., Adolphs, R., Hurley, R., & Piven, J. (2007). Abnormal use of facial information in high-functioning autism. *Journal of Autism and Developmental Disorders*, *37*, 929–939.
- Tantam, D., Monaghan, L., Nicholson, H., & Stirling, J. (1989). Autistic children's ability to interpret faces: A research note. *Journal of Child Psychology and Psychiatry*, *30*, 623–630.
- Teunisse, J. P., & de Gelder, B. (2001). Impaired categorical perception of facial expressions in high-functioning adolescents with autism. *Child Neuropsychology (Neuropsychology, Development and Cognition: Section C)* *7*, 1–14.
- Thomas, M. S. C., Annaz, D., Ansari, D., Scerif, G., Jarrold, C., & Karmiloff-Smith, A. (2009). Using developmental trajectories to understand developmental disorders. *Journal of Speech Language and Hearing Research*, *52*, 336–358.
- Tracy, J., Robins, R., Schriber, R., & Solomon, M. (2011). Is emotion recognition impaired in individuals with autism spectrum disorders? *Journal of Autism and Developmental Disorders*, *41*, 102–109.
- Van den Noortgate, W., & Onghena, P. (2006). Analysing repeated measures data in cognitive research: A comment on regression coefficient analyses. *The European Journal of Cognitive Psychology*, *18*, 937–952.
- Van Der Geest, J. N., Kemner, C., Verbaten, M. N., & Van Engeland, H. (2002). Gaze behavior of children with pervasive developmental disorder toward human faces: A fixation time study. *Journal of Child Psychology and Psychiatry*, *43*, 669–678.
- Walker-Andrews, A. S. (1998). Emotions and social development: Infants' recognition of emotions in others. *Pediatrics*, *102*, e1268.
- Wallace, S., Coleman, M., & Bailey, A. (2008). An investigation of basic facial expression recognition in autism spectrum disorders. *Cognition and Emotion*, *22*, 1353–1380.
- Weeks, S. J., & Hobson, R. P. (1987). The salience of facial expression for autistic children. *Journal of Child Psychology and Psychiatry*, *28*, 137–152.
- Whalen, P. J. (2007). The uncertainty of it all. *Trends in Cognitive Sciences*, *11*, 499–500.

Table 1: Studies of emotion recognition in ASC. * indicates datasets included in the primary analysis; - indicates information that was not available; EL = emotion labelling task, EM = emotion matching task, EFAS = Ekman facial affect set

Row	Study	Autism group				Control Group				Task	Results			
		number (males)	Diagnosis (tools)	Mean age in Years (SD)	Level of function	number (males)	Mean age in Years (SD)	Level of function	type		stimuli	Emotions	mean ASD (SD)	Mean control (SD)
1	*Atkinson, 2009	13 (12)	autism spectrum disorder	30.9 (13.8)	FSIQ: 116.2 (12.2) VIQ: 106.9 (11.6) PIQ: 105.2 (13.5)	16	26.7 (12.8)	FSIQ: 106.6 (8.5) VIQ: 105.7 (10.0) PIQ: 108.4 (4.8)	EL (forced)	Patch-light and full-light displays	Anger, disgust, fear, happiness, sadness	0.61 (0.195)	0.846 (0.196)	-1.1664 (0.1629)
2	*Baron-Cohen, Spitz & Cross, 1993	15	autism (Rutter)	12.6 (3.5)	VMA: 5.3	15	4.4 (0.3)	-	EM	photos / drawings from books	happy, sad, surprised	7.7 (2.18)	8.5 (1.54)	-0.4124 (0.1362)
3	*Bolte & Poustka, 2003	15 (12)	autism (german ADOS)	15.7 (8.6)	NVIQ: 103.7 (23.7)	22 (11)	29.7 (10.3)	NVIQ: 112.9 (8.4)	EL (forced)	Ekman facial affect set + author pictures	6 basic + neutral	30.07 (8.2)	42.9 (13)	-1.0538 (0.1271)
4	*Boraston et al., 2007 experiment 2	11 (9)	ASD ADOS	36.7	VIQ: 118 (118-130) PIQ: 117 (107-133)	9(7)	34 (15)	VIQ: 107 (15.6) PIQ: 112 (12.5)	EL (forced)	EFAS	6 basic	7.64 (2.10)	8.69 (0.96)	-0.5923 (0.2108)
5a	*Braverman et al., 1989	10	autism	10.9 (2.7)	DAD MA: 6.2 (2.3)	10	5.4 (1.3)	DAD: 6.2 (1.8)	EL (forced)	Ekman facial affect set	happy, sad, mad, scared	13 (2.66)	15.10 (1.28)	-0.9636 (0.2232)
5b	Braverman et al., 1989	10	autism	10.9 (2.7)	DAD MA: 6.2 (2.3)	10	5.4 (1.3)	DAD: 6.2 (1.8)	EM	Ekman facial affect set	happy, sad, mad, scared	9 (2.94)	13.30 (3.20)	-1.3403 (0.2449)
5c	Braverman et al., 1989	10	autism	10.9 (2.7)	DAD MA: 6.2 (2.3)	10	5.4 (1.3)	DAD: 6.2 (1.8)	AC	Ekman facial affect set	happy, sad, mad, scared	10.50 (2.99)	14.30 (2.16)	-1.3954 (0.2487)
6a	*Buitelaar et al., 1999	20	autism	12.5 (3.2)	FSIQ: 102.1 (19.0) VIQ :104.1 (15.4) PIQ: 99.5 (21.8)	20	10.5 (1.9)	-	EM	Ekman facial affect set	happiness, sadness, anger, fear	12.8 (2.95)	14.45 (1.43)	-0.6976 (0.1061)
6b	Buitelaar et al., 1999	20	autism	12.5 (3.2)	FSIQ: 102.1 (19.0) VIQ :104.1 (15.4) PIQ: 99.5 (21.8)	20	10.5 (1.9)	-	EM (context)	Ekman facial affect set	happiness, sadness, anger, fear	6.05 (1.09)	6.45 (0.60)	-0.4456 (0.1025)
7	*Castelli, 2005 experiment 2	20	autism, Asperger syndrome	12.3 (2.3)	Verbal MA: 9.2 (2.6) Performance: 10.1 (3.2)	20	9,2 (2.4)	Verbal MA: 9.11 (2.7)	EL (free naming)	Ekman facial affect set	6 basic + neutral	7.5 (2.6)	6.9 (2.8)	0.2177 (0.1006)
8	*Celani, Battacchi & Arcidiacono, 1999	10 (8)	autism	12.7 (3.8)	VMA: 7.0 (2.1) IQ: 63.2 (19.4)	10 (8)	6.3 (1.6)	VMA: 6.6 (1.3) IQ: 101.6 (6.6)	EM (delayed)	Ekman facial affect set + Ladavas, 1982	Happy, sad, "wry faces" (distractors)	4.60 (1.26)	7 (0.47)	-2.4172 (0.3461)
9	*Clark, Winkelman & McIntosh, 2008	15 (13)	ASD ADI-R, ADOS	26 (7.5)	PPVT: 99.5 (23.8)	10(9)	19.7 (4.6)	PPVT: 112.9 (4.8)	EL (forced)	Ekman facial affect set	happy / angry	0.59 (0.16)	0.76 (0.09)	-1.2005 (0.1955)

Row	Study	Autism group				Control Group				Task		Results		
		number (males)	Diagnosis (tools)	Mean age in Years (SD)	Level of function	number (males)	Mean age in Years (SD)	Level of function	type	stimuli	Emotions	mean ASD (SD)	Mean control (SD)	Hedge's d (Var d)
10	*Corden, Chilvers & Skuse, 2008	21 (16)	Asperger syndrome ADI-R, ADOS	33.8 (13.6)	FSIQ: 117.9 (11.67) VIQ: 115.8 (9.68) Non-VIQ: 116.2 (13.73)	21 (16)	32.1 (11.58)	FSIQ: 117.2 (8.00) VIQ: 115.1 (8.19) Non-VIQ: 115.5 (8.75)	EL (forced)	EFAS	6 basic	8.11 (1.69)	8.77 (1.37)	-0.4175 (0.0973)
11	*Da Fonseca et al, 2009	19 (17)	autism, Asperger syndrome ADI-R	12.8 (3.6)	IQ: 93.7 (15.4)	19 (17)	12.7 (3.4)	-	EM (context)	Visual scene with masked faces	fear, sadness, anger, happiness	68.7 (2.32)	81.3 (2.33)	-5.3056 (0.4757)
12a	*Davies et al., 1994 Experiment 2	10	autism	13.72 (1.91)	BPVS MA: 5.65 (1.62) Raven's Matrices MA: 8.0 (1.59)	20	13.63 (1.56)	BPVS MA: 6.3 (2.05) Raven's Matrices MA: 7.13 (2.99)	EM	Ekman facial affect set	,happy, sad, surprised, angry	18.4 (4.81)	15.5 (5.23)	0.5534 (0.1551)
12b	*Davies et al., 1994 Experiment 2	9	autism, Asperger syndrome	14.35 (1.49)	Mill Hill MA: 12.2 (2.3) Raven's Matrices MA: 10.72 (3.16)	11	13.93 (1.45)	Mill Hill MA: 11.5 (1.69) Raven's matrices: 9.72 (1.76)	EM	Ekman facial affect set	,happy, sad, surprised, angry	20 (1.32)	21.1 (1.57)	-0.7195 (0.215)
13a	*Deruelle, Rondan, Gepner & Tardiff, 2004	11	ASD (CARS)	6.7 (2.4)	IQ: 80	11	6.6 (2.3)	-	EM	Black-and-white photographs of 25 adult faces and five faces of children	Disgust, surprise and happiness	65.7 (16.8)	90.3 (10.3)	-1.6984 (0.2474)
13b	Deruelle, Rondan, Gepner & Tardiff, 2004	11	ASD (CARS)	6.7 (2.4)	IQ: 80	11	6.6 (2.3)	-	EM	Black-and-white photographs of 25 adult faces and five faces of children	Disgust, surprise and happiness	65.7 (16.8)	96.1 (5.1)	-2.3557 (0.3079)
14	*Downs & Smith, 2004	10	autism ADI-R	7.1 (1.1)	IQ: 106 (7.63)	10	7.7 (1.2)	108.9 (12.22)	-	Level 1 (identifying emotional facial)	-	3.3 (1.6)	4 (0)	-0.5926 (0.2088)
15a	*Dyck, Ferguson & Shochet, 2001	28 (24)	Asperger syndrome	-	-	36 (27)	-	-	EL (free naming)	Facial cues test	6 basic + contempt + neutral	18.92 (4.12)	20.27 (3.12)	-0.3715 (0.0646)
15b	*Dyck, Ferguson & Shochet, 2001	20 (17)	autism	-	-	36 (27)	-	-	EL (free naming)	Facial cues test	6 basic + contempt + neutral	12.15 (5.9)	20.27 (3.12)	-1.7785 (0.106)
16	*Dziobek et al., 2006	17 (14)	Asperger syndrome ADI-R, ASDI	41.4 (9.9)	Shipley IQ: 113 (6)	17 (15)	40.2 (13)	Shipley IQ: 115 (5)	EL (forced)	Ekman facial affect set	6 basic + neutral	21.6 (2.9)	25.4 (1.9)	-1.5134 (0.1513)

Row	Study	Autism group				Control Group				Task	Results			
		number (males)	Diagnosis (tools)	Mean age in Years (SD)	Level of function	number (males)	Mean age in Years (SD)	Level of function	type		stimuli	Emotions	mean ASD (SD)	Mean control (SD)
17a	*Fein et al., 1992	15	PDD	128.53 (30.82) months	DAD MA: 73.6 (27.07) PPVT: 63.6 (15.22)	15	60.2 (11.87) months	PPVT matched: 62.67 (14.89) DAD MA: 63.57 (14.74)	EM (context)	Pictures of affect laden context (face of child was not visible)	happy, sad, angry, scared	9.67 (3.62)	10.87 (2.75)	-0.3632 (0.1355)
17b	Fein et al., 1992	15	PDD	128.53 (30.82) months	DAD MA: 73.6 (27.07) PPVT: 63.6 (15.22)	15	64.73 (18.70) months	DAD MA matched: 74 (26.99) PPVT: 75.27 (27.92)	EM (context)	Pictures of affect laden context (face of child was not visible)	happy, sad, angry, scared	9.67 (3.62)	11.07 (4.25)	-0.3451 (0.1353)
18a	Gepner, Dereulle & Grynfeldt, 2001	13	autism	69.38 (11) months	Developmental age (French Brunet-Lezine scale, 1951): 40.53 (13.6) months	13	40.7 (13.6)	Matched on developmental age	EM	Videotaped sequences of female's face displaying expressions. Strob face condition	Joy, surprise, sadness, disgust	2.46 (1.36)	3 (1.09)	-0.4243 (0.1573)
18b	Gepner, Dereulle & Grynfeldt, 2001	13	autism	69.38 (11) months	Developmental age (French Brunet-Lezine scale, 1951): 40.53 (13.6) months	13	40.7 (13.6)	Matched on developmental age	EM	Videotaped sequences of female's face displaying expressions. Dynamic face condition	Joy, surprise, sadness, disgust	2.86 (1.51)	2.69 (1.49)	0.1098 (0.1541)
18c	*Gepner, Dereulle & Grynfeldt, 2001	13	autism	69.38 (11) months	Developmental age (French Brunet-Lezine scale, 1951): 40.53 (13.6) months	13	40.7 (13.6)	Matched on developmental age	EM	Videotaped sequences of female's face displaying expressions. Still face condition	Joy, surprise, sadness, disgust	2.61 (1.38)	2.38 (1.44)	0.1579 (0.1543)
19	*Gepner, de Gelder & de Schonen 1996 (task 3b)	7 (4)	autism	11.13 (4.8)	Verbal test: 125 (40.2) Non verbal test: 19.41 (5.84)	7	5.11 (1.10)	Nonverbal test: 19 (5.54)	EM	Black and white photographs from Campbell, Landis & Regard, 1986	happy, surprise, dislike, neutral	32.1 (6.69)	80.4 (2.6)	-8.9093 (3.1206)
19	Gepner, de Gelder & de Schonen 1996 (task 3b)	7 (4)	autism	11.13 (4.8)	Verbal test: 125 (40.2) Non verbal test: 19.41 (5.84)	7	5.7 (2.5)	Verbal test: 131.5 (30.9)	EM	Black and white photographs from Campbell, Landis & Regard, 1986	happy, surprise, dislike, neutral	32.1 (6.69)	75 (4.04)	-7.3353 (2.2074)
20	*Grossman & Tager-Flusberg, 2008	25 (22)	autism ADI-R, ADOS	13.8 (3.3)	VIQ: 105.04 (19.35) NonVIQ: 107.56(10.80) PPVT: 111.63 (19.97)	25 (20)	14.1 (3.0)	VIQ: 112.60 (14.58) NonVIQ: 113 (12.57) PPVT: 115.29 (9.6)	Aranging pictures in timeline	Professional actress portraying emotions, 6 still images were extracted from each clip	Happy, sad, anger, fear, disgust	46.33 (21.12)	61 (18.59)	-0.7258 (0.0853)

Row	Study	Autism group				Control Group				Task		Results		
		number (males)	Diagnosis (tools)	Mean age in Years (SD)	Level of function	number (males)	Mean age in Years (SD)	Level of function	type	stimuli	Emotions	mean ASD (SD)	Mean control (SD)	Hedge's d (Var d)
21	*Grossman et al., 2000	13 (12)	Asperger syndrome	11.8 (3.27)	FSIQ: 106.4 (18.423) VIQ: 115.8 (15.627) PIQ: 97.2 (21.586)	13 (12)	11.5 (1.898)	Matched on FSIQ and VIQ not on PIQ $t(1,24)=2.039$, $p<0.10$	EL (forced)	Ekman facial affect set	Happy, sad, angry, afraid, surprised	0.750 (0.296)	0.827 (0.234)	-0.2898 (0.1555)
22	*Hadjikhani et al., 2009	11 (9)	autism spectrum disorder ADI-R, ADOS	30 (11)	IQ: 126 (10)	11 (8)	31 (14)	-	EM	Bodily expressions of emotions with actor's face obscured (upright condition)	Sadness, anger, fear	15.5 (2.7)	17.3 (1.0)	-0.8505 (0.1983)
23	*Hobson, 1986	23 (18)	autism Rutter (1974)	14.7 (30.4 mo)	Raven's matrices: 29.9 (13.5)	23 (15)	7.3 (21 mo)	Raven's matrices: 29.4 (12.1)	EM	Video clips showing bodily expressions of emotions with actor's face obscured	Anger, sadness, happiness, fear	5.69 (3.262)	9.782 (2.569)	-1.3695 (0.1073)
24	*Hubert et al., 2006	16 (14)	autism spectrum disorder ASSQ scale	25.6 (9.2)	IQ: 92.2	16 (14)	27.2 (10.6)	-	EL (forced)	Video sequences displaying faces of 8 actors expressing emotion	Happiness / anger	97.9 (10.8)	99.8 (9.1)	-0.1855 (0.1255)
25	*Hubert et al., 2007	19 (17)	autism, Asperger syndrome ADI-R, ASSQ scale	21.6 (6.1)	IQ: 83.3 (15.9)	19 (17)	24.4 (8.6)	-	EL (free, descriptive)	Point-light displays	Surprised, angry, frightened, sad, happy	31.42 (25.6)	77.14 (15.4)	-2.1189 (0.1643)
26	*Humphreys et al., 2007 experiment 1	20	autism ADI-R, ADOS	24 (9)	FSIQ: 103 (15) VIQ: 102 (14) PIQ: 105 (15)	18	28 (10)	FSIQ: 109 (9) VIQ: 107 (11) PIQ: 108 (7)	EL (forced)	FEEST (Young et al., 2002) set of morphed facial expressions	6 basic	81.19 (24.25)	94.24 (11.76)	-0.6592 (0.1113)
27	*Jones et al, 2010	97	ADOS	15.5(5.6)	VIQ: 81.1(17.9) PIQ: 90.6 (18.6)	55	15.5 (5.9)	VIQ: 86.3 (20.2) PIQ: 91.5 (21.7)	EL(forced)	Ekman facial affect set	6 basic	6.99 (2.19)	7.28 (1.96)	-0.1367 (0.0286)
28	*Kircher, Hatri, Heekeren, & Dziobek, 2011	15 (5)	ASD, ADI-R	31.9 (7.6)	IQ: 112.6 (11.6)	27	31.8 (7.4)	IQ: 110.1(8.7)	EL	Multifaceted Empathy Test (MET) (Dziobek et al. 2008)	-	94 (11.9)	96.8 (4.6)	-0.3452 (0.1051)
29a	*Lacroix et al., 2007	12 (6)	autism ADI-R, ADOS	6.1 (1.2)	VMA: 5.8 (1.11)	12 (6)	5.7 (1.10)	VMA: 5.7 (1.10)	EL (free)	Facial expressions posed according to standard facial configuration	Happy, sad, angry, afraid, surprised, neutral	0.73 (0.52)	0.65 (0.43)	0.1619 (0.1672)

		Autism group				Control Group			Task			Results		
Row	Study	number (males)	Diagnosis (tools)	Mean age in Years (SD)	Level of function	number (males)	Mean age in Years (SD)	Level of function	type	stimuli	Emotions	mean ASD (SD)	Mean control (SD)	Hedge's d (Var d)
29b	Lacroix et al., 2007	12 (6)	autism ADI-R, ADOS	6.1 (1.2)	VMA: 5.8 (1.11)	12 (6)	5.7 (1.10)	VMA: 5.7 (1.10)	EM	Facial expressions posed according to standard facial configuration	Happy, sad, angry, afraid, surprised, neutral	0.71 (0.19)	0.75 (0.26)	-0.1696 (0.1673)
29c	Lacroix et al., 2007	12 (6)	autism ADI-R, ADOS	6.1 (1.2)	VMA: 5.8 (1.11)	12 (6)	5.7 (1.10)	VMA: 5.7 (1.10)	EL (forced)	Facial expressions posed according to standard facial configuration	Happy, sad, angry, afraid, surprised, neutral	0.8 (0.19)	0.88 (0.18)	-0.4174 (0.1703)
30a	*Linder & Rosen, 2006	14 (12)	Asperger syndrome	10.21 (2.89)	PPVT-III: 114.86 (17.15)	16 (11)	10.19 (3.12)	PPVT-III: 117.31 (14.30)	EM	Perception of Emotion Test- (Egan, 1989) Static facial expression	Happy, angry, sad, neutral	12.57 (2.62)	14.38 (1.20)	-0.8851 (0.147)
30b	Linder & Rosen, 2006	14 (12)	Asperger syndrome ASDS scale, PDD checklist	10.21 (2.89)	PPVT-III: 114.86 (17.15)	16 (11)	10.19 (3.12)	PPVT-III: 117.31 (14.30)	EM	Perception of Emotion Test- (Egan, 1989) Dynamic facial expression	Happy, angry, sad, neutral	14.21 (1.67)	15.30 (0.81)	-0.8265 (0.1453)
31a	*Macdonald et al., 1989	10 (10)	autism	27.2 (5.6)	BPVS MA: 84.1 (27.6) Raven's matrices IQ: 118.4 (13)	10 (10)	26.2 (2.9)	Raven's matrices IQ: 120.1 (8.6)	EL (free)	Ekman facial affect et	Happiness, sadness, anger, fear, neutral	16.2 (3.0)	19.6 (0.6)	-1.5052 (0.2566)
31b	Macdonald et al., 1989	10 (10)	autism	27.2 (5.6)	BPVS MA: 84.1 (27.6) Raven's matrices IQ: 118.4 (13)	10 (10)	26.2 (2.9)	Raven's matrices IQ: 120.1 (8.6)	EM (context)	Pictures of affect laden context (face of child was not visible)	Happiness, sadness, anger, fear, neutral	15.1 (3.9)	19.2 (0.6)	-1.4074 (0.2495)
32	*O'Connor, 2007	18	Asperger syndrome	26.9 (7.8)	-	18	25.2 (6.5)	-	EL (forced)	Facial expressions from Mind Reading Emotions Library (Baron-cohen et al., 2003)	Happy, sad, angry	92.93 (8.86)	95.53 (5.64)	-0.3423 (0.1127)
33a	*Ozonoff, Pennington & Rogers, 1990	13 (9)	autism CARS	6.24 (2.14)	Leiter IQ: 76 (20) Non-verbal MA: 4.57 (1.43)	13 (9)	4.14 (1.49)	Leiter IQ: 114 (20) Non-verbal MA: 4.56 (1.44)	EM (faces)	Same facial expressions	Happiness, sadness, anger	6.31 (2.10)	8.92 (2.29)	-1.1504 (0.1793)
33b	Ozonoff, Pennington & Rogers, 1990	14 (10)	autism CARS	6.40 (2.04)	Verbal MA: 3.38 (0.39)	14 (10)	3 (0.27)	Verbal MA: 3.54 (0.51)	EM (faces)	Same facial expressions	Happiness, sadness, anger	6.57 (2.24)	6.79 (2.22)	-0.0958 (0.143)
33c	Ozonoff, Pennington & Rogers, 1990	13 (9)	autism CARS	6.24 (2.14)	Leiter IQ: 76 (20) Non-verbal MA: 4.57 (1.43)	13 (9)	4.14 (1.49)	Leiter IQ: 114 (20) Non-verbal MA: 4.56 (1.44)	ES	Facial expressions posed according to standard facial configuration	Happiness / sadness	23.31 (4.31)	27 (1.92)	-1.0711 (0.1759)

Row	Study	Autism group				Control Group				Task			Results	
		number (males)	Diagnosis (tools)	Mean age in Years (SD)	Level of function	number (males)	Mean age in Years (SD)	Level of function	type	stimuli	Emotions	mean ASD (SD)	Mean control (SD)	Hedge's d (Var d)
33d	Ozonoff, Pennington & Rogers, 1990	14 (10)	autism CARS	6.40 (2.04)	Verbal MA: 3.38 (0.39)	14 (10)	3 (0.27)	Verbal MA: 3.54 (0.51)	ES	Facial expressions posed according to standard facial configuration	Happiness / sadness	24 (4.17)	26 (2.18)	-0.5836 (0.1489)
33e	Ozonoff, Pennington & Rogers, 1990	13 (9)	autism CARS	6.24 (2.14)	Leiter IQ: 76 (20) Non-verbal MA: 4.57 (1.43)	13 (9)	4.14 (1.49)	Leiter IQ: 114 (20) Non-verbal MA: 4.56 (1.44)	EM (context)	Pictures of affect laden context (face of child was not visible)	Happiness, sadness, amger	4.92 (2.33)	8.31 (2.50)	-1.3586 (0.1893)
33f	Ozonoff, Pennington & Rogers, 1990	14 (10)	autism CARS	6.40 (2.04)	Verbal MA: 3.38 (0.39)	14 (10)	3 (0.27)	Verbal MA: 3.54 (0.51)	EM (context)	Pictures of affect laden context (face of child was not visible)	Happiness, sadness, amger	5.29 (2.61)	5.86 (1.75)	-0.2491 (0.144)
34	*Pelphrey et al., 2002	5 (5)	autism ADI-R, ADOS	25.2	FSIQ: 100.75 (7.69) VIQ: 117 (23.12)	5 (5)	28.20	-	EL (forced)	Ekman facial affect set	6 basic	76 (11.53)	92.60 (8.11)	-1.5042 (0.5131)
35a	*Philip et al., 2009	23 (16)	autism spectrum disorder ADI-R	32.5 (10.9)	FSIQ: 101.5 (18.5) VIQ: 98.2 (15.8) PIQ: 104.4 (18.6)	23 (17)	32.4 (11.1)	FSIQ: 111.2 (8.5) VIQ: 106.8 (8.8) PIQ: 113.4 (10.4)	EL (forced)	Ekman facial affect set	6 basic	70.74 (14.79)	88.30 (5.47)	-1.5478 (0.113)
35b	Philip et al., 2009	23 (16)	autism spectrum disorder ADI-R	32.5 (10.9)	FSIQ: 101.5 (18.5) VIQ: 98.2 (15.8) PIQ: 104.4 (18.6)	23 (17)	32.4 (11.1)	FSIQ: 111.2 (8.5) VIQ: 106.8 (8.8) PIQ: 113.4 (10.4)	EL (forced)	JACFEE series (Matsumoto and Ekman, 1988)	Happiness, anger, fear, disgust, sadness	79.61 (16.62)	91.83 (9.57)	-0.8857 (0.0955)
35c	Philip et al., 2009	23 (16)	autism spectrum disorder ADI-R	32.5 (10.9)	FSIQ: 101.5 (18.5) VIQ: 98.2 (15.8) PIQ: 104.4 (18.6)	23 (17)	32.4 (11.1)	FSIQ: 111.2 (8.5) VIQ: 106.8 (8.8) PIQ: 113.4 (10.4)	EL (forced)	Video displaying emotions with whole body movement (face not visible)	Happiness, anger, fear, disgust, sadness	71.48 (15.28)	86.17 (7.18)	-1.2094 (0.1029)
35d	Philip et al., 2009	23 (16)	autism spectrum disorder ADI-R	32.5 (10.9)	FSIQ: 101.5 (18.5) VIQ: 98.2 (15.8) PIQ: 104.4 (18.6)	23 (17)	32.4 (11.1)	FSIQ: 111.2 (8.5) VIQ: 106.8 (8.8) PIQ: 113.4 (10.4)	EM	JACFEE series	Happiness, anger, fear, disgust, sadness	83.61 (10.87)	93.09 (9.48)	-0.9136 (0.096)
36a	*Piggot et al., 2004	14 (14)	autism ADI-R, ADOS-G	13.1 (2.5)	FSIQ: 112 (15.9); VIQ: 104 (20.3); PIQ: 118 (13.6)	14 (14)	14.4 (3.3)	FSIQ: 116 (10.5); VIQ: 114 (14.2); PIQ: 114 (6.3)	EL	Ekman facial affect set	Fearful, surprised, angry	69 (27)	85 (12)	-0.7435 (0.1527)
36b	Piggot et al., 2004	14 (14)	autism ADI-R, ADOS-G	13.1 (2.5)	FSIQ: 112 (15.9); VIQ: 104 (20.3); PIQ: 118 (13.6)	14 (14)	14.4 (3.3)	FSIQ: 116 (10.5); VIQ: 114 (14.2); PIQ: 114 (6.3)	EM	Ekman facial affect set	Fearful, surprised, angry	76 (25)	90 (11)	-0.7038 (0.1517)
37	*Robel et al., 2004	14	autism	8.44 (1.74)	VIQ: 88 (16.2); PIQ: 95 (9.4)	20	7.9 (1.31)	-	EM	The Minnesota Test of Affective Processing	-	0.869 (0.065)	0.865 (0.076)	0 (0.1214)

Autism group						Control Group			Task			Results		
Row	Study	number (males)	Diagnosis (tools)	Mean age in Years (SD)	Level of function	number (males)	Mean age in Years (SD)	Level of function	type	stimuli	Emotions	mean ASD (SD)	Mean control (SD)	Hedge's d (Var d)
38a	*Rosset et al., 2008	20 (19)	autism spectrum disorder ADI-R	9.4 (3.2)	MA: 8.2 (4.0)	20 (19)	-	MA matched: 8 (4)	EL (forced)	Photographs of human face, human cartoon faces, non-human cartoon faces	Sadness, happiness, anger	2.7 (2.5)	3.4 (3.4)	-0.2299 (0.1007)
38b	Rosset et al., 2008	20 (19)	autism spectrum disorder ADI-R	9.4 (3.2)	MA: 8.2 (4.0)	20 (19)	9.6 (3.3)	matched on CA and gender	EL (forced)	Photographs of human face, human cartoon faces, non-human cartoon faces	Sadness, happiness, anger	2.7 (2.5)	2.5 (2.6)	0.0769 (0.1001)
39	*Rump et al., 2009 experiment 1	19 (14)	autism ADOS-G	6.4 (0.8)	VMA: 97.89 (15.86)	18 (11)	6 (0.8)	105.61 (11.99)	EL (free)	Dynamic face stimuli created by authors	Happy, sad, angry, afraid	1.80 (0.52)	2.42 (0.59)	-1.0928 (0.1243)
40	*Rutherford & Towns, 2008	11 (11)	autism spectrum disorder ADOS	25.8 (6.09)	FSIQ: 91.8 (14.07) VIQ: 97.1 (18.28) PIQ: 86.7 (12.12)	11 (11)	25.7 (8.87)	FSIQ: 101.1 (12.40) VIQ: 101.1 (11.77) PIQ: 99.3 (15.03)	EL (forced)	Face photos from Baron-Cohen et al. (1997)	6 basic + distress	82 (12.8)	88 (6)	-0.5774 (0.1894)
41	*Sawyer, Williamson & Young, 2011	48	AS	21.6 (9.8)	FSIQ: 108 (17.9) VIQ: 109 (19.2) PIQ: 104 (18.2)	27	24 (9.2)	FSIQ: 114 (13) VIQ: 113 (12.8) PIQ: 111.4 (12.8)	EL	Colour photographs of actors posing based on Ekman faces	happy, sad, angry, afraid, surprised, disgust	78.6 (15.8)	94.8 (7.7)	-1.189 (0.0673)
42	*Spezio et al., 2007	9 (9)	autism ADI-R, ADOS	23	FSIQ: 107 VIQ: 109 PIQ: 104	10 (10)	28	FIQ: 106 VIQ: 101 PIQ: 111	EL (forced)	Ekamn facial affect set, "Bubbles" task	Happy, fearful	82 (3)	80 (5)	0.4571 (0.2166)
43a	*Tantam et al., 1989	10 (8)	autism	12.14 (3.10)	Raven's 22.3 (7.8) PIQ: 67.3 (20.4) EPVT 29.1 (11) VIQ: 42.3 (9.6)	10 (9)	12.19 (3.13)	Raven's 21.2 (6.9) PIQ: 62.3 (17.5) EPVT 46.4 (17.8) VIQ: 56.5 (12.7)	EL (forced) (just test 4)	Ekman facial affect set	6 basic	7.6 (4.5)	12.2 (4.7)	-0.9575 (0.2229)
43b	Tantam et al., 1989	10 (8)	autism	12.14 (3.10)	Raven's 22.3 (7.8) PIQ: 67.3 (20.4) EPVT 29.1 (11) VIQ: 42.3 (9.6)	10 (9)	12.19 (3.13)	Raven's 21.2 (6.9) PIQ: 62.3 (17.5) EPVT 46.4 (17.8) VIQ: 56.5 (12.7)	Odd one out	Ekman facial affect set	6 basic	12.5 (5.9)	23.7 (4.6)	-2.0277 (0.3028)
44	*Tracy, Robins, Schriber, & Solomon, 2011	29	ASD	11.7 (2.6)	FSIQ: 110 (21.5) VIQ: 111 (19.6) PIQ: 110 (18.9)	31	12.3 (2.5)	FSIQ: 119 (12.5) VIQ: 115 (15.7) PIQ: 113 (11.8)	EL	UCDavis Set of Emotion Expressions (UCDSSEE; Tracy et al.2009)	6 basic + pride + contempt	79 (23.7)	76 (27.4)	0.1153 (0.0669)

Row	Study	Autism group				Control Group			Task			Results		
		number (males)	Diagnosis (tools)	Mean age in Years (SD)	Level of function	number (males)	Mean age in Years (SD)	Level of function	type	stimuli	Emotions	mean ASD (SD)	Mean control (SD)	Hedge's d (Var d)
45	*Wallace et al., 2011	42 (38)	11 HFA, 27AS, 3 PDD-NOS, 1 ASD	15.7 (2.79)	FSIQ: 113.80 (16.11) VIQ: 112.18 (16.36) PIQ: 110.40 (14.93)	31 (28)	16.35(2.0)	FSIQ: 113.74 (10.94) VIQ: 111.16 (11.79) PIQ: 112.74 (10.47)	EL	The Emotional MultiMorph Task	6 basic	82.39 (15.88)	91.94 (9.33)	-0.6996 (0.0594)
46	*Wallace, Coleman & Bailey, 2008, expt 1	26(26)	autism/asperger (clinician), ADI-R	31(9)	BPVS raw: 148(13) Raven's IQ 101(18)	26(23)	32(9)	BPVS raw: 153(9) Raven's IQ 98(12)	EL(forced)	EFAS	6 basic + neutral	70.83 (20.53)	85.83 (16.18)	-0.7991 (0.0831)
47	*Wicker et al., 2008	12 (11)	autism spectrum disorder	27 (11)	FSIQ :81.83 VIQ: 83.17 NVIQ: 81.58	14	23.4 (10)	Matched on age not on IQ	EL (forced)	Dynamic facial expressions	Angry, happy	91.28 (10.8)	96.51 (9.1)	-0.5108 (0.1598)
48a	*Wright et al., 2008	35 (33)	autism, Asperger syndrome ADI-R, ADOS	11.31 (2.17)	FSIQ: 104.63 (17.99) VIQ: 105.66 (21.01) PIQ: 103.03 (16.09)	35 (33)	11.57 (1.94)	FSIQ: 103.86 (16.26) VIQ: 105.74 (16.31) PIQ: 100.94 (16.39)	EL (forced)	Ekman facial affect set	6 basic	40.26 (9.56)	41.51 (7.99)	-0.1403 (0.0573)
48b	Wright et al., 2008	33	autism, Asperger syndrome ADI-R, ADOS	-	-	33	-	-	EL (forced)	Pictures with visual contextual information in which Individual showed facial expression	Fear, disgust, anger, surprise, sadness	39.8 (5.53)	39.9 (5.09)	-0.0186 (0.0606)

row no	Study	number autistic	age autistic	number control	age control	Task	happy	sad	angry	surprised	fearful	disgusted
1a	Atkinson, 2009	13	30.9	16	26.7	EL	-0.79	-0.41	-1.04		-0.76	-0.89
1b	Atkinson, 2009	13	30.9	16	26.7	EL	-1.09		-1.41		-0.67	-1.15
2	Baron-Cohen, Spitz & Cross, 1993	15	12.6	15	4.4	EM	-0.06	0.17		-1.08		
3	Boraston et al., 2007 expt2b	11	36.7	9	34	EL	-0.41	-1.31	-0.57		-0.52	
4	Castelli, 2005 expt2	20	12.3	20	9.2	EL	0.00	0.68	0.21	0.08	-0.04	0.46
5	Corden, Chilvers & Skuse, 2008	21	33.8	21	32.1	EL	-0.58	-0.40	-0.16	-0.16	-0.93	-0.61
6	Grossman et al., 2000	13	11.8	13	11.5	EL	-0.55	-0.27	-0.56	-0.53	-0.07	
7	Humphreys et al., 2007 expt 1 (unambiguous)	20	24	18	28	EL	-0.41	-0.56	-0.93	-0.54	-1.34	-0.36
8	Jones et al	97	15.5	55	15.5	EL	0.04	0.09	-0.32	-0.28	0.04	-0.30
9a	Lacroix et al., 2007	12	6.1	12	5.7	EL	0.62	0.00	-0.14	-0.16	0.06	
9b	Lacroix et al., 2007	12	6.1	12	5.7	EM	-0.18	-0.88	0.16	0.38	0.17	
9c	Lacroix et al., 2007	12	6.1	12	5.7	EL	-0.34	-1.04	-0.71	-0.28	-0.17	
10	Linder & Rosen, 2006	14	10.21	16	10.19	EM	-0.40	-0.61	-0.75			
11	O'Connor, 2007	18	26.9	18	25.2	EL	-0.33	-0.34	-0.40			
12	Pelphrey et al., 2002	5	25.2	5	28.2	EL	-0.57	-0.53	-1.21	-0.46	-1.53	-0.33
13a	Philip et al., 2009	23	32.5	23	32.4	EM	0.00	-0.94	-0.67		-0.56	-0.64
13b	Philip et al., 2009	23	32.5	23	32.4	EL	-0.03	-0.56	-0.98		-0.17	-0.69
13c	Philip et al., 2009	23	32.5	23	32.4	EL	-1.23	-0.47	-0.70		-1.18	-0.88
14	Tracy, Robins, Schriber, & Solomon, 2011	29	11.7	31	12.3	EL	-0.37	-0.08	0.29	-0.32	0.06	-0.31
15	Wallace et al., 2011	42	15.7	31	16.34	EL	0.27	-0.41	-0.81	-0.24	-0.47	-0.32
16	Wallace, Coleman & Bailey, 2008 just upright faces	26	32	26	31	EL	-0.57	-1.02	-0.55	-0.72	-1.09	-0.86
	Mean		21.24		20.27		-0.22	-0.33	-0.45	-0.37	-0.46	-0.40
	upper confidence limit						0.00	-0.10	-0.21	-0.10	-0.20	-0.12
	lower confidence limit						-0.44	-0.55	-0.68	-0.65	-0.71	-0.69
	Number included	379		329			16	16	15	11	13	10

Paper	Number of participants with ASD	brief description of results	Reason for exclusion
Adolphs, Sears, & Piven (2001)	7	mostly no difference	means / sd not available.
Ashwin, Wheelwright, & Baron-Cohen (2006)	13 (expt 1) 26 (expt 2)	ASD worse than TD	means / sd not available.
Balconi, & Carrera (2007)	7	mixed results	means / sd not available.
Baron-Cohen et al. (1997)	16	both groups at ceiling	means / sd not available.
Bauminger (2004)	16	subtle differences	emotion type / task
Begeer, Terwogt, Rieffe, Stegge, Olthof, & Koot (2010)	31	subtle differences	stimuli type / task
Begeer, Banerjee, Rieffe, Terwogt, Potharst, Stegge, & Koot (2011)	22 (study 1) 25 (study 2)	na	stimuli type / task
Bormann-Kischkel, Vilsmeier & Baude (1995)	41	ASD worse than TD	means / sd not available.
Buitelaar, & van der Weesn (1997)	20 AD; 20 PDD-NOS	mixed results	stimuli type / task means / sd not available
Critchley, Daly, Bullmore, Williams, Van Amelsvoort, Robertson et al. (2000)	9	ASD worse than TD	stimuli type / task means / sd not available
Dalton, Holsen, Abbeduto, & Davidson (2008)	14	ASD worse than TD	stimuli type / task
Dalton, Nacewicz, Johnstone, Schaefer, Gernsbacher, Goldsmith, Alexander, & Davidson (2008)	14	ASD worse than TD	stimuli type / task
Dawson, Webb, Carver, Panagiotides, & McPartland J (2004)	29	significant difference in N300 to fearful vs neutral faces	stimuli type / task
Dennis, Lockyer, & Lazenby (2000)	8	ASD worse than TD	emotion type / task
Golan, Baron-Cohen, & Golan (2008)	23	ASD worse than TD	emotion type / task
Golan, Baron-Cohen, Hill, & Golan (2006)	22	ASD worse than TD	emotion type / task
Gross (2004)	27	ASD worse than TD	stimuli type / task control groups means / sd not available
Gross (2005)	24	ASD worse than TD	stimuli type / task control groups means / sd not available
Harris, & Lindell (2011)	127	na	sample
Heerey, Keltner, & Capps (2003)	25	no group differences	means / sd not available.
Hertzog, Snow, & Sherman (1989)	18	na	stimuli type / task
Howard et al. 2000	10	ASD only worse at fear	means / sd not available.
Joseph, Ehrman, McNally, & Keehn (2008)	20	subtle differences	stimuli type / task means / sd not available
Kamio, Wolf, & Fein (2006)	16	na	stimuli type / task
Kamio, & Toichi (1998)			not able to find the full article
Kätstyri, Saalasti, Tiippana, von Wendt, & Sams (2008)	20	no differences	stimuli type / task
Kikuchi, & Koga (2001)			not able to find the full article
Krysko, Kristen M, & Rutherford (2009)	19	ASD worse than TD	stimuli type / task
Kuusikko, Haapsamo, Jansson-Verkasalo, Hurtig, Mattila, Ebeling, Jussila, Bölte, & Moilanen (2009)	57 (26 HFA; 31 AS)	ASD worse than TD	means / sd not available.
Laine, Rauzy, Tardif, & Gepner (2011)	19		stimuli type / task
Losh et al. (2009)	36	mixed results	means / sd not available.
Loveland et al. (2008)	80	ASD worse than TD	means / sd not available.
Loveland, TunaliKotoski, Chen, et al. (1997)	35	no overall group differences	means / sd not available.
McHugh, Bobarnac, & Reed (2010)	3		sample size/no control group
Miyahara, Bray, Tsujii, Fujita, & Sugiyama (2007)	20	no overall group differences	stimuli type / task
Moore, Hobson, & Lee (1997)	13	ASD worse than TD	stimuli type / task
Njiokiktjien, Vershoor, de Sonnevile, Huyser, Op het Veld, & Toorenaar (2001)	3	emotion recognition deficits in all three participants	sample size/no control group
Nueman, Spezio, Piven & Adolphs (2006)	10	no group differences	means / sd not available.
O'Connor, Hamm & Kirk, 2005	15	mixed results	means / sd not available.
Ozonoff, Pennington, & Rogers (1991)	23	no significant difference	emotion type
Ozonoff, Rogers, & Pennington (1991)	13	ASD worse than TD	emotion type
Prior, Dahlstrom, & Squires (1990)	20	no group differences	means / sd not available.
Riefe, Tervogt, & Kotronopoulou (2007)	22	ASD worse than TD at recognition of own emotions	stimuli type / task
Rieffe, Meerum Terwogt, & Stockmann(2000)	23	subtle differences	stimuli type / task
Rose, Lincoln, Lai, Ene, Searcy, & Bellugi (2007)	16	ASD worse than Williams syndrome for upright faces, opposite pattern for inverted faces	stimuli type / task
Rutherford, & McIntosh (2007)	10	ASD different to TD	stimuli type / task
Ryan, & Charragáin, (2010)	20	significant improvements in emotional recognition after training	no control group
Shamay-Tsoory, Tomer, Yaniv, & Aharon-Peretz (2002)	2	no significant impairment in emotion recognition	sample size/no control group

Paper	Number of participants with ASD	brief description of results	Reason for exclusion
Shamay-Tsoory (2008)	18	ASD worse than TD	emotion type
Tardiff et al. (2007)	12	ASD worse than TD	means / sd not available.
Teunisse, & de Gelder (2001)	17	more impairments in ASD individuals with lower social intelligence	means / sd not available.
van der Geest, Kemner, Verbaten, & van Engeland (2002)	17	no differences in gaze patterns	stimuli type / task
Volkmar, Sparrow, Rende, & Cohen (1989)	16	no differences in face perception	stimuli type / task
Wang, Dapretto, Hariri, Sigman, & Bookheimer (2004)	12	subtle differences	stimuli type / task
Wang, Lee, Sigman, & Dapretto (2007)	18	no group differences	stimuli type / task
Wicker, Fonlupt, Hubert, Tardif, Gepner, & Deruelle (2008)	12	no group differences	stimuli type / task
Wong et al. 2008	10	no group differences	means / sd not available.
Yirmiya, Sigman, Kasari, & Mundy (1992)	18	ASD worse than TD	stimuli type / task

Full References of all studies in the meta-analysis

Studies in Table 1.

Atkinson, A. P. (2009). Impaired recognition of emotions from body movements is associated with elevated motion coherence thresholds in autism spectrum disorders. *Neuropsychologia*, *47*(13), 3023-3029.

Baron-cohen, S., Spitz, A., & Cross, P. (1993). Do children with autism recognise surprise? A research note. *Cognition & Emotion*, *7*(6), 507-516.

Bolte, S., & Poustka, F. (2003). The recognition of facial affect in autistic and schizophrenic subjects and their first-degree relatives. *Psychological medicine*, *33*(5), 907-915.

Boraston, Z., Blakemore, S. J., Chilvers, R., & Skuse, D. (2007). Impaired sadness recognition is linked to social interaction deficit in autism. *Neuropsychologia*, *45*(7), 1501-1510.

Braverman, M., Fein, D., Lucci, D., & Waterhouse, L. (1989). Affect comprehension in children with pervasive developmental disorders. *Journal of autism and developmental disorders*, *19*(2), 301-316.

Buitelaar, J. K., Van der Wees, M., Swaab-Barneveld, H. A. N. N. A., & Van Der Gaag, R. J. (1999). Theory of mind and emotion-recognition functioning in autistic spectrum disorders and in psychiatric control and normal children. *Development and Psychopathology*, *11*(1), 39-58.

Castelli, F. (2005). Understanding emotions from standardized facial expressions in autism and normal development. *Autism*, *9*(4), 428-449.

Celani, G., Battacchi, M. W., & Arcidiacono, L. (1999). The understanding of the emotional meaning of facial expressions in people with autism. *Journal of autism and developmental disorders*, *29*(1), 57-66.

Clark, T. F., Winkielman, P., & McIntosh, D. N. (2008). Autism and the extraction of emotion from briefly presented facial expressions: Stumbling at the first step of empathy. *Emotion*, *8*(6), 803.

Corden, B., Chilvers, R., & Skuse, D. (2008). Avoidance of emotionally arousing stimuli predicts social-perceptual impairment in Asperger's syndrome. *Neuropsychologia*, *46*(1), 137-147.

Da Fonseca, D., Santos, A., Bastard-Rosset, D., Rondan, C., Poinso, F., & Deruelle, C. (2009). Can children with autistic spectrum disorders extract emotions out of contextual cues?. *Research in Autism Spectrum Disorders*, *3*(1), 50-56.

Davies, S., Bishop, D., Manstead, A. S., & Tantam, D. (2006). Face perception in children with autism and Asperger's syndrome. *Journal of Child Psychology and Psychiatry*, *35*(6), 1033-1057.

Deruelle, C., Rondan, C., Gepner, B., & Tardif, C. (2004). Spatial frequency and face processing in children with autism and Asperger syndrome. *Journal of autism and developmental disorders*, *34*(2), 199-210.

Downs, A., & Smith, T. (2004). Emotional understanding, cooperation, and social behavior in high-functioning children with autism. *Journal of autism and developmental disorders*, *34*(6), 625-635.

Dyck, M. J., Ferguson, K., & Shochet, I. M. (2001). Do autism spectrum disorders differ from each other and from non-spectrum disorders on emotion recognition tests?. *European child & adolescent psychiatry*, *10*(2), 105-116.

Dziobek, I., Fleck, S., Rogers, K., Wolf, O. T., & Convit, A. (2006). The 'amygdala theory of autism' revisited: Linking structure to behavior. *Neuropsychologia*, *44*(10), 1891-1899.

- Fein, D., Lueci, D., Braverman, M., & Waterhouse, L. (1992). Comprehension of affect in context in children with pervasive developmental disorders. *Journal of Child Psychology and Psychiatry*, 33(7), 1157-1162.
- Gepner, B., Deruelle, C., & Grynfeldt, S. (2001). Motion and emotion: A novel approach to the study of face processing by young autistic children. *Journal of autism and developmental disorders*, 31(1), 37-45.
- Gepner, B., de Gelder, B., & de Schonen, S. (1996). Face processing in autistics: Evidence for a generalised deficit?. *Child Neuropsychology*, 2(2), 123-139.
- Grossman, R. B., & Tager-Flusberg, H. (2008). Reading faces for information about words and emotions in adolescents with autism. *Research in autism spectrum disorders*, 2(4), 681-695.
- Grossman, J. B., Klin, A., Carter, A. S., & Volkmar, F. R. (2000). Verbal bias in recognition of facial emotions in children with Asperger syndrome. *Journal of Child Psychology and Psychiatry*, 41(3), 369-379.
- Hadjikhani, N., Joseph, R. M., Manoach, D. S., Naik, P., Snyder, J., Dominick, K., ... & De Gelder, B. (2009). Body expressions of emotion do not trigger fear contagion in autism spectrum disorder. *Social cognitive and affective neuroscience*, 4(1), 70-78.
- Hobson, R. P. (1986). The autistic child's appraisal of expressions of emotion. *Journal of Child Psychology and Psychiatry*, 27(3), 321-342.
- Hubert, B., Wicker, B., Moore, D. G., Monfardini, E., Duverger, H., Fonseca, D. D., & Deruelle, C. (2007). Brief report: recognition of emotional and non-emotional biological motion in individuals with autistic spectrum disorders. *Journal of autism and developmental disorders*, 37(7), 1386-1392.
- Humphreys, K., Minshew, N., Leonard, G. L., & Behrmann, M. (2007). A fine-grained analysis of facial expression processing in high-functioning adults with autism. *Neuropsychologia*, 45(4), 685-695.
- Jones, C. R., Pickles, A., Falcato, M., Marsden, A. J., Happé, F., Scott, S. K., ... & Charman, T. (2011). A multimodal approach to emotion recognition ability in autism spectrum disorders. *Journal of Child Psychology and Psychiatry*, 52(3), 275-285.
- Kirchner, J. C., Hatri, A., Heekeren, H. R., & Dziobek, I. (2011). Autistic symptomatology, face processing abilities, and eye fixation patterns. *Journal of autism and developmental disorders*, 41(2), 158-167.
- Lacroix, A., Guidetti, M., & Reilly, J. (2009). Recognition of emotional and nonemotional facial expressions: A comparison between Williams syndrome and autism. *Research in developmental disabilities*.
- Lindner, J. L., & Rosén, L. A. (2006). Decoding of emotion through facial expression, prosody and verbal content in children and adolescents with Asperger's syndrome. *Journal of Autism and Developmental Disorders*, 36(6), 769-777.
- Macdonald, H., Rutter, M., Howlin, P., Rios, P., Conteur, A. L., Evered, C., & Folstein, S. (1989). Recognition and expression of emotional cues by autistic and normal adults. *Journal of Child Psychology and Psychiatry*, 30(6), 865-877.
- O'Connor, K. (2007). Brief report: Impaired identification of discrepancies between expressive faces and voices in adults with Asperger's syndrome. *Journal of autism and developmental disorders*, 37(10), 2008-2013.

- Ozonoff, S., Pennington, B. F., & Rogers, S. J. (1990). Are there emotion perception deficits in young autistic children?. *Journal of Child Psychology and Psychiatry*, 31(3), 343-361.
- Pelphrey, K. A., Sasson, N. J., Reznick, J. S., Paul, G., Goldman, B. D., & Piven, J. (2002). Visual scanning of faces in autism. *Journal of autism and developmental disorders*, 32(4), 249-261.
- Philip, R. C. M., Whalley, H. C., Stanfield, A. C., Sprengelmeyer, R., Santos, I. M., Young, A. W., ... & Hall, J. (2010). Deficits in facial, body movement and vocal emotional processing in autism spectrum disorders. *Psychological medicine*, 40(11), 1919-1929.
- Piggot, J., Kwon, H., Mobbs, D., Blasey, C., Lotspeich, L., Menon, V., ... & Reiss, A. L. (2004). Emotional attribution in high-functioning individuals with autistic spectrum disorder: a functional imaging study. *Journal of the American Academy of Child & Adolescent Psychiatry*, 43(4), 473-480.
- Robel, L., Ennouri, K., Piana, H., Vaivre-Douret, L., Perier, A., Flament, M. F., & Mouren-Siméoni, M. C. (2004). Discrimination of face identities and expressions in children with autism: Same or different?. *European child & adolescent psychiatry*, 13(4), 227-233.
- Rosset, D. B., Rondan, C., Da Fonseca, D., Santos, A., Assouline, B., & Deruelle, C. (2008). Typical emotion processing for cartoon but not for real faces in children with autistic spectrum disorders. *Journal of autism and developmental disorders*, 38(5), 919-925.
- Rump, K. M., Giovannelli, J. L., Minshew, N. J., & Strauss, M. S. (2009). The development of emotion recognition in individuals with autism. *Child development*, 80(5), 1434-1447.
- Rutherford, M. D., & Towns, A. M. (2008). Scan path differences and similarities during emotion perception in those with and without autism spectrum disorders. *Journal of autism and developmental disorders*, 38(7), 1371-1381.
- Sawyer, A. C., Williamson, P., & Young, R. L. (2012). Can Gaze Avoidance Explain Why Individuals with Asperger's Syndrome Can't Recognise Emotions From Facial Expressions?. *Journal of autism and developmental disorders*, 42(4), 606-618.
- Spezio, M. L., Adolphs, R., Hurley, R. S., & Piven, J. (2007). Abnormal use of facial information in high-functioning autism. *Journal of autism and developmental disorders*, 37(5), 929-939.
- Tantam, D., Monaghan, L., Nicholson, H., & Stirling, J. (2006). Autistic children's ability to interpret faces: a research note. *Journal of Child Psychology and Psychiatry*, 30(4), 623-630.
- Tracy, J. L., Robins, R. W., Schriber, R. A., & Solomon, M. (2011). Is emotion recognition impaired in individuals with autism spectrum disorders?. *Journal of autism and developmental disorders*, 41(1), 102-109.
- Wallace, G. L., Case, L. K., Harms, M. B., Silvers, J. A., Kenworthy, L., & Martin, A. (2011). Diminished sensitivity to sad facial expressions in high functioning autism spectrum disorders is associated with symptomatology and adaptive functioning. *Journal of autism and developmental disorders*, 41(11), 1475-1486.
- Wicker, B., Fonlupt, P., Hubert, B., Tardif, C., Gepner, B., & Deruelle, C. (2008). Abnormal cerebral effective connectivity during explicit emotional processing in adults with autism spectrum disorder. *Social cognitive and affective neuroscience*, 3(2), 135-143.
- Wright, B., Clarke, N., Jordan, J. O., Young, A. W., Clarke, P., Miles, J., ... & Williams, C. (2008). Emotion recognition in faces and the use of visual context Vo in young people with high-functioning autism spectrum disorders. *Autism*, 12(6), 607-626.

Studies in Table 2

These studies are all included in the list above for Table 1.

Studies in Table 3

Adolphs, R., Sears, L., & Piven, J. (2001). Abnormal processing of social information from faces in autism. *Journal of Cognitive neuroscience*, 13(2), 232-240.

Ashwin, C., Chapman, E., Colle, L., & Baron-Cohen, S. (2006). Impaired recognition of negative basic emotions in autism: A test of the amygdala theory. *Social Neuroscience*, 1(3-4), 349-363.

Balconi, M., & Carrera, A. (2007). Emotional representation in facial expression and script: A comparison between normal and autistic children. *Research in developmental disabilities*, 28(4), 409-422.

Baron-Cohen, S., Jolliffe, T., Mortimore, C., & Robertson, M. (1997). Another advanced test of theory of mind: Evidence from very high functioning adults with autism or Asperger syndrome. *Journal of Child Psychology and Psychiatry*, 38(7), 813-822.

Bauminger, N. (2004). The expression and understanding of jealousy in children with autism. *Development and psychopathology*, 16(1), 157-177.

Begeer, S., Terwogt, M. M., Rieffe, C., Stegge, H., Olthof, T., & Koot, H. M. (2010). Understanding emotional transfer in children with autism spectrum disorders. *autism*, 14(6), 629-640.

Bormann-Kischkel, C., Vilsmeier, M., & Baude, B. (1995). The development of emotional concepts in autism. *Journal of child psychology and psychiatry*, 36(7), 1243-1259.

Buitelaar, J. K., & van der Wees, M. (1997). Are deficits in the decoding of affective cues and in mentalizing abilities independent?. *Journal of autism and developmental disorders*, 27(5), 539-556.

Critchley, H. D., Daly, E. M., Bullmore, E. T., Williams, S. C., Van Amelsvoort, T., Robertson, D. M., ... & Murphy, D. G. (2000). The functional neuroanatomy of social behaviour changes in cerebral blood flow when people with autistic disorder process facial expressions. *Brain*, 123(11), 2203-2212.

Dalton, K. M., Holsen, L., Abbeduto, L., & Davidson, R. J. (2008). Brain function and gaze fixation during facial-emotion processing in fragile X and autism. *Autism Research*, 1(4), 231-239.

Dalton, K. M., Nacewicz, B. M., Johnstone, T., Schaefer, H. S., Gernsbacher, M. A., Goldsmith, H. H., ... & Davidson, R. J. (2005). Gaze fixation and the neural circuitry of face processing in autism. *Nature neuroscience*, 8(4), 519-526.

Dawson, G., Webb, S. J., Carver, L., Panagiotides, H., & McPartland, J. (2004). Young children with autism show atypical brain responses to fearful versus neutral facial expressions of emotion. *Developmental science*, 7(3), 340-359.

Dennis, M., Lockyer, L., & Lazenby, A. L. (2000). How high-functioning children with autism understand real and deceptive emotion. *Autism*, 4(4), 370-381.

Golan, O., Baron-Cohen, S., & Golan, Y. (2008). The 'Reading the Mind in Films' task [child version]: Complex emotion and mental state recognition in children with and without autism spectrum conditions. *Journal of Autism and Developmental Disorders*, 38(8), 1534-1541.

- Golan, O., Baron-Cohen, S., & Hill, J. (2006). The Cambridge Mindreading (CAM) Face-Voice Battery: Testing complex emotion recognition in adults with and without Asperger syndrome. *Journal of autism and developmental disorders*, 36(2), 169-183.
- Gross, T. F. (2004). The perception of four basic emotions in human and nonhuman faces by children with autism and other developmental disabilities. *Journal of Abnormal Child Psychology*, 32(5), 469-480.
- Gross, T. F. (2005). Global–local precedence in the perception of facial age and emotional expression by children with autism and other developmental disabilities. *Journal of autism and developmental disorders*, 35(6), 773-785.
- Harris, C. D., & Lindell, A. K. (2011). The influence of autism-like traits on cheek biases for the expression and perception of happiness. *Brain and cognition*, 77(1), 11-16.
- Heerey, E. A., Keltner, D., & Capps, L. M. (2003). Making sense of self-conscious emotion: linking theory of mind and emotion in children with autism. *Emotion*, 3(4), 394.
- Hertzig, M. E., Snow, M. E., & Sherman, M. (1989). Affect and cognition in autism. *Journal of the American Academy of Child & Adolescent Psychiatry*, 28(2), 195-199.
- Howard, M. A., Cowell, P. E., Boucher, J., Broks, P., Mayes, A., Farrant, A., & Roberts, N. (2000). Convergent neuroanatomical and behavioural evidence of an amygdala hypothesis of autism. *Neuroreport*, 11(13), 2931-2935.
- Joseph, R. M., Ehrman, K., McNally, R., & Keehn, B. (2008). Affective response to eye contact and face recognition ability in children with ASD. *Journal of the International Neuropsychological Society*, 14(6), 947.
- Kamio, Y., Wolf, J., & Fein, D. (2006). Automatic processing of emotional faces in high-functioning pervasive developmental disorders: An affective priming study. *Journal of autism and developmental disorders*, 36(2), 155-167.
- Kätsyri, J., Saalasti, S., Tiippana, K., von Wendt, L., & Sams, M. (2008). Impaired recognition of facial emotions from low-spatial frequencies in Asperger syndrome. *Neuropsychologia*, 46(7), 1888-1897.
- Krysko, K. M., & Rutherford, M. D. (2009). A threat-detection advantage in those with autism spectrum disorders. *Brain and cognition*, 69(3), 472-480.
- Kuusikko, S., Haapsamo, H., Jansson-Verkasalo, E., Hurtig, T., Mattila, M. L., Ebeling, H., ... & Moilanen, I. (2009). Emotion recognition in children and adolescents with autism spectrum disorders. *Journal of autism and developmental disorders*, 39(6), 938-945.
- Lainé, F., Rauzy, S., Tardif, C., & Gepner, B. (2011). Slowing Down the Presentation of Facial and Body Movements Enhances Imitation Performance in Children with Severe Autism. *Journal of autism and developmental disorders*, 41(8), 983-996.
- Losh, M., Adolphs, R., Poe, M. D., Couture, S., Penn, D., Baranek, G. T., & Piven, J. (2009). Neuropsychological profile of autism and the broad autism phenotype. *Archives of general psychiatry*, 66(5), 518.
- Loveland, K. A., Bachevalier, J., Pearson, D. A., & Lane, D. M. (2008). Fronto-limbic functioning in children and adolescents with and without autism. *Neuropsychologia*, 46(1), 49-62.

- McHugh, L., Bobarnac, A., & Reed, P. (2011). Brief Report: Teaching Situation-Based Emotions to Children with Autistic Spectrum Disorder. *Journal of autism and developmental disorders*, 41(10), 1423-1428.
- Miyahara, M., Bray, A., Tsujii, M., Fujita, C., & Sugiyama, T. (2007). Reaction time of facial affect recognition in Asperger's disorder for cartoon and real, static and moving faces. *Child Psychiatry & Human Development*, 38(2), 121-134.
- Moore, D. G., Hobson, R. P., & Lee, A. (2011). Components of person perception: An investigation with autistic, non-autistic retarded and typically developing children and adolescents. *British Journal of Developmental Psychology*, 15(4), 401-423.
- Njiokiktjien, C., Verschoor, A., De Sonnevile, L., Huyser, C., Op het Veld, V., & Toorenaar, N. (2001). Disordered recognition of facial identity and emotions in three Asperger type autists. *European child & adolescent psychiatry*, 10(1), 79-90.
- Neumann, D., Spezio, M. L., Piven, J., & Adolphs, R. (2006). Looking you in the mouth: abnormal gaze in autism resulting from impaired top-down modulation of visual attention. *Social Cognitive and Affective Neuroscience*, 1(3), 194-202.
- O'Connor, K., Hamm, J. P., & Kirk, I. J. (2005). The neurophysiological correlates of face processing in adults and children with Asperger's syndrome. *Brain and Cognition*, 59(1), 82-95.
- Ozonoff, S., Pennington, B. F., & Rogers, S. J. (1991). Executive function deficits in high-functioning autistic individuals: relationship to theory of mind. *Journal of child psychology and psychiatry*, 32(7), 1081-1105.
- Ozonoff, S., Rogers, S. J., & Pennington, B. F. (1991). Asperger's Syndrome: Evidence of an Empirical Distinction from High-Functioning Autism. *Journal of Child Psychology and Psychiatry*, 32(7), 1107-1122.
- Prior, M., Dahlstrom, B., & Squires, T. L. (1990). Autistic children's knowledge of thinking and feeling states in other people. *Journal of Child Psychology and Psychiatry*, 31(4), 587-601.
- Rieffe, C., Meerum Terwogt, M., & Kotronopoulou, K. (2007). Awareness of single and multiple emotions in high-functioning children with autism. *Journal of autism and developmental disorders*, 37(3), 455-465.
- Rieffe, C., Terwogt, M. M., & Stockmann, L. (2000). Understanding atypical emotions among children with autism. *Journal of Autism and Developmental Disorders*, 30(3), 195-203.
- Rose, F. E., Lincoln, A. J., Lai, Z., Ene, M., Searcy, Y. M., & Bellugi, U. (2007). Orientation and affective expression effects on face recognition in Williams syndrome and autism. *Journal of autism and developmental disorders*, 37(3), 513-522.
- Rutherford, M. D., & McIntosh, D. N. (2007). Rules versus prototype matching: Strategies of perception of emotional facial expressions in the autism spectrum. *Journal of autism and developmental disorders*, 37(2), 187-196.
- Ryan, C., & Charragáin, C. N. (2010). Teaching emotion recognition skills to children with autism. *Journal of autism and developmental disorders*, 40(12), 1505-1511.
- Shamay-Tsoory, S. G., Tomer, R., Yaniv, S., & Aharon-Peretz, J. (2002). Empathy deficits in Asperger syndrome: a cognitive profile. *Neurocase*, 8(3), 245-252.

- Shamay-Tsoory, S. G. (2008). Recognition of 'fortune of others' emotions in Asperger syndrome and high functioning autism. *Journal of autism and developmental disorders*, 38(8), 1451-1461.
- Tardif, C., Lainé, F., Rodriguez, M., & Gepner, B. (2007). Slowing down presentation of facial movements and vocal sounds enhances facial expression recognition and induces facial-vocal imitation in children with autism. *Journal of autism and developmental disorders*, 37(8), 1469-1484.
- Teunisse, J. P., & de Gelder, B. (2001). Impaired categorical perception of facial expressions in high-functioning adolescents with autism. *Child Neuropsychology*, 7(1), 1-14.
- Van der Geest, J. N., Kemner, C., Verbaten, M. N., & Van Engeland, H. (2002). Gaze behavior of children with pervasive developmental disorder toward human faces: a fixation time study. *Journal of Child Psychology and Psychiatry*, 43(5), 669-678.
- Volkmar, F. R., Sparrow, S. S., Rende, R. D., & Cohen, D. J. (1989). Facial perception in autism. *Journal of Child Psychology and Psychiatry*, 30(4), 591-598.
- Wang, A. T., Dapretto, M., Hariri, A. R., Sigman, M., & Bookheimer, S. Y. (2004). Neural correlates of facial affect processing in children and adolescents with autism spectrum disorder. *Journal of the American Academy of Child & Adolescent Psychiatry*.
- Wang, A. T., Lee, S. S., Sigman, M., & Dapretto, M. (2006). Neural basis of irony comprehension in children with autism: the role of prosody and context. *Brain*, 129(4), 932-943.
- Wicker, B., Fonlupt, P., Hubert, B., Tardif, C., Gepner, B., & Deruelle, C. (2008). Abnormal cerebral effective connectivity during explicit emotional processing in adults with autism spectrum disorder. *Social cognitive and affective neuroscience*, 3(2), 135-143.
- Wong, T. K., Fung, P. C., Chua, S. E., & McAlonan, G. M. (2008). Abnormal spatiotemporal processing of emotional facial expressions in childhood autism: dipole source analysis of event-related potentials. *European journal of neuroscience*, 28(2), 407-416.
- Yirmiya, N., Sigman, M. D., Kasari, C., & Mundy, P. (1992). Empathy and Cognition in High-Functioning Children with Autism. *Child development*, 63(1), 150-160.